

The three Millennium problem solutions, RH, NSE, YME, and a Hilbert scale based quantum geometrodynamics

Klaus Braun
September 19, 2020

www.fuchs-braun.com

„looking back, part (A)“

on a 10 year journey to ...

Riemann Hypothesis solutions

A modified Zeta function theory is proposed to overcome current challenges

- (a) to verify several **Riemann Hypothesis** (RH) criteria
- (b) to prove the binary **Goldbach conjecture**

The current two baseline functions to define the Zeta functions, the Gaussian function and the (periodical) fractional part function (resp. their corresponding Mellin transforms) are replaced by their corresponding Hilbert transforms, which are the Dawson function (which is a specific Kummer function) and the Fourier series representation of the $\log(\sin x)$ -function. The convergence analysis is based on corresponding Hilbert space frameworks, supporting especially Cardon's „convolution operator representation“ and Bagchi's "Nyman-Beurling" RH criteria applied to the modified entire Zeta function. Thereby, the convolution operator representation goes along with convergent (Mellin transform) integrals, overcoming the corresponding challenge of an only formally valid self-adjoint invariant operator representation of the standard entire Zeta function ((EdH) 10.3).

Basically, the Bagchi RH criterion and the Cardon RH criterion are two sides of the same coin, which is about the construction of appropriately defined operators, i.e. a defined mapping rule in combination with a defined domain.

The Bagchi-"Nyman-Beurling" RH criteria comes along with the $\log(\sin x) L_2^\#(0,1)$ -function (the Hilbert transform of the fractional part function $\rho(x)$) in the form

$$\rho_H(x) = \sum_1^{\infty} \frac{\cos 2\pi vx}{\pi v} = -\frac{1}{\pi} \log 2 \sin(\pi x) \in L_2^\#(0,1)$$

defined in the (periodical) Hilbert scale framework $H_0^\#(0,1)$. It enables the definition of new arithmetical functions going along with a Hilbert space based "circle method" defined on the „boundary of the unit circle“, alternatively to the "open unit disk" domain of the Hardy-Littlewood circle method. The counterpart of the zeros of the e^{ix} -function are the zeros of the Kummer function ${}_1F_1\left(\frac{1}{2}; \frac{3}{2}; -x\right) = \sum_{n=0}^{\infty} \frac{(-1)^n x^n}{2^{n+1} n!}$ enabling the definition of a pair of two different arithmetical functions to analyse (p, q) -binary number theoretical problems. The concept also supports the verification of the Snirelmann density criterion to prove the Goldbach conjecture.

The proposed modified Zeta function theory supports the proof of several RH criteria, which might be grouped into the following three classes (A1)-(A3), basically defined by the applied underlying function space frameworks:

(A1) this class is about RH criteria which can be re-formulated in terms of distributional Hilbert scale functions $H_\alpha(-\infty, \infty)$

(A2) this class is about RH criteria which can be re-formulated in terms of periodical distributional Hilbert scale functions $H_\alpha^\#(0,1)$

(A3) this class is about RH criteria formulated by distributional arithmetical functions going beyond the distributional arithmetical functions applied for "a distributional way to prove the Prime Number Theorem" (ViJ).

The common key challenge is in the context of the "Polya theorem", where the not-vanishing constant Fourier term provides the key handicap. From the related "Polya theorem" pdf (see pdf in the "literature" page) we quote:

"Attempts have been made to apply Polya's general theorem about the zeros of the Fourier transform of a real function to the Zeta function (PoG). After a change of variable $t \rightarrow \log x$ and approximating the integral over the half-line (positive x -axis) by integrals over finite intervals (which are the essential restriction/handicap of Polya's theorem to be applied to the Zeta function), one can obtain a theorem about zeros of Mellin transforms. For the Zeta function, the Müntz formula has been used, but the Polya theorem only applies in the interval $(0,1)$, and no information is obtained about the zeros of the Zeta function in the critical open stripe of validity of Müntz's formula".

(OIJ): „The theory of Dirichlet series offers a bridge between number theory and analysis. Perhaps the most appealing example of the power of this connection is given by the tauberian approach to the classical prime number theorem.“ ... The asymptotic behavior in terms of the related Chebyshev-type inequality can be connected to local function theoretic properties of (distributional) (Bagchi-type) Hilbert spaces H_ω .

The Hilbert-Polya conjecture (which is about the existence of a proper self-adjoint integral operator going along with the concept of convolution operators, (CaD)) needs to overcome the mathematical problem of the not vanishing constant Fourier term of the Jacobian theta function.

Every Hilbert transformed L_2 function has a vanishing constant Fourier term. The Paley-Wiener functions form a Hilbert subspace of PW of $L_2(\mathbb{R})$, which is a reproducing kernel Hilbert space (HiJ). Thus, the inner product of any $f \in PW$ with $w := w(t, x) = \frac{\sin(\pi(t-x))}{\pi(t-x)}$, i.e. $f(t) = (f, w)_0 = \int_{-\infty}^{\infty} f(x)w(t-x)dt$. PW and PW^\perp are invariant subspaces for the Hilbert transforms on $L_2(\mathbb{R})$. Calderón's reproducing formula provided the baseline for the continuous wavelet transformation theory coming along with the concept of „windowed Fourier transforms“ (LoA). The PW_π reproducing kernel provided the baseline for the cardinal series theory, addressing e.g. the question „how power series coefficients a_n of a function $f(z) = \sum a_n z^n$ determine its singularities“, (HiJ). Corresponding π -cardinal series are convergent under the condition $\sum \frac{|a_n|}{n} < \infty$. Cardinal series can be obtained formally by considering the Lagrange interpolation formula, which is proposed alternatively to the Euler-Mclaurin /Newton-Gauss summation formulas.

The canonical resolution of the „ E_μ spectrum of the dynamical Hamiltonian mechanics system“ (consisting of a discontinuous and a continuous part) is concerned with certain self-adjoint and unitary operators in a Hilbert space. The conception that such a spectrum reveals the mechanical properties of the system in its own structure lead to the concepts of an inner product $(E_\mu f, g)$ on the Hilbert space, PW -like formulas (KoB1), which is the well known spectral theory in Hilbert spaces for unitary operators (e.g. (HiF)). The physical motivations for related ergodic theorems are based on the Maxwell-Boltzmann gas theory and the Gibbs statistical mechanics (HoE).

With respect to part B, the common mathematical tools to analyze the proposed complementary kinematical energy space H_1^\perp and the non-linear Landau damping phenomenon are

- the wavelets, where a vanishing constant Fourier term of a L_2 function is a sufficient condition to be a wavelet function (HoM), (MeY); the underlying „theory of frames“, especially the Nonharmonic, but exact Fourier frames are closely related to the Riesz basis of a separable Hilbert space, (YoR) p. 157, (NaA); the latter one plays a key role in the context of the Kadec 1/4-theorem (related to the Paley-Wiener space PW (DuR), resp. indefinite metrics, (AzT), (BoJ), as applied in (A2), (A3) below; general references to frames are provided in (YoR) p. 190.
- ergodic theory on a Hilbert space (HaP), which is about measure-preserving transformation, e.g. in relationship to turbulence problems in hydrodynamics (HoE), (LiP), in combination with non-harmonic Fourier series theory (YoR), based on complex exponentials forming a Riesz basis for $L_2^\#$ and sequences of real or complex numbers with uniform density one (DuR).

The parts (A1-A3) below are concerned with Hilbert scales in the form $H_\alpha, l_2^\alpha, \alpha \in \mathbb{R}$:

The Dirichlet series theory is an extension of the concept of power series replacing $\sum_1^\infty a_n e^{-xn} \rightarrow \sum_1^\infty a_n e^{-x \log n}$. The relationship between the Dirichlet series

$$f(s) := \sum_1^\infty a_n e^{-s \log n} \quad g(s) := \sum_1^\infty b_n e^{-s \log n}$$

and the Hilbert space $H_{-1/2}^\# \cong l_2^{-1/2}$ on the critical line is given by ([LaE] §227, Satz 40):

$$((f, g))_{-1/2} := \lim_{\omega \rightarrow \infty} \frac{1}{2\omega} \int_{-\omega}^{\omega} f\left(\frac{1}{2} + it\right) g(1/2 - it) dt = \sum_1^\infty \frac{1}{n} a_n b_n.$$

The cardinal series theory is an extension of the Dirichlet series theory. In our case this is about absolute convergent series $C(x) = \frac{\sin(\pi(x-\lambda))}{\pi} \sum_{n=1}^\infty (-1)^n \frac{c(n+\lambda)}{x-\lambda-n}$ with cardinal series $C(x) =$

$\frac{w}{\pi} \sum_{-\infty}^\infty a_n \frac{\sin \frac{w}{\pi}(x-a-nw)}{x-a-nw}$ for the points $(a+nw, a_n)$, whereby $\sum_2^\infty (|a_n| + |a_{-n}|) \frac{\log n}{n} < \infty$ to ensure that the cardinal series $\frac{\sin \pi x}{\pi} \left[\frac{a_0}{x} + \sum_{n=1}^\infty (-1)^n \left(\frac{a_n}{x-n} + \frac{a_{-n}}{x+n} \right) \right]$ is absolutely convergent (WhJ1).

Remark: The inverse mapping of $\sigma(x) := \tilde{T}(\log x) := \sum_{n \leq x} \frac{1}{n} \log \left(\frac{x}{n} \right)$, $x \geq 1$, is given by ((ScW), lemma 3.3, p.36) $\sigma^{-1}(x) = \tilde{T}^{-1}(\log x) = \sum_{n \leq x} \frac{\mu(n)}{n} \log \left(\frac{x}{n} \right)$. The link to the Riemann, von Mangoldt and Landau density functions $J(x) := \frac{1}{2} \left[\sum_{p^n < x} \frac{1}{n} + \sum_{p^n \leq x} \frac{1}{n} \right]$, $\psi(x) := \sum_{n \leq x} \Lambda(n)$, $\vartheta(x) = \sum_{n=1}^\infty \Lambda(n) \log \left(\frac{x}{n} \right) = \psi(x) \log x$ (fulfilling $d[x\vartheta'] = d\psi = \log x dj$) is given by $x\sigma'(x) = \sum_{n \leq x} \frac{1}{n}$. The considered sequences below enable modified Riemann density functions $J^*(x), \psi^*(x), \vartheta^*(x)$ with $dJ = dJ^*, d\psi^* = d\psi, d\vartheta = d\vartheta^*$ (see also lemma 5 v), vii), viii) below).

From (ApT) p. 66 we recall that the PNT can be derived from the formula $\sum_{n=1}^\infty \frac{\mu(n)}{n} = 0$. What cannot be derived from the PNT is the convergence of the series, (LaE) §160, $\sum_{n=1}^\infty \frac{\mu(n)}{n} \log \left(\frac{1}{n} \right) = -\sum_{n=2}^\infty \mu(n) \frac{\log n}{n} = 1$.

Putting $a_n := \sqrt{n} \mu(n) \log \left(\frac{1}{n} \right)$ ($a_n^* := \mu(n) \log \left(\frac{1}{n} \right)$) and $b_n := \frac{1}{\sqrt{n}}$ ($b_n^* := 1$) one gets

$$((f, g))_{-1/2} = ((f^*, g^*))_{-1/2} = 1 \quad \text{and} \quad ((g, g))_{-1/2} = ((g^*, g^*))_{-1} = \sum_1^\infty \frac{1}{n} b_n^2 = \sum_1^\infty \frac{1}{n^2} = \frac{\pi^2}{6},$$

i.e. $f\left(\frac{1}{2} + it\right), g\left(\frac{1}{2} + it\right) \in H_{-1/2}$ resp. $g^*\left(\frac{1}{2} + it\right) \in H_{-1}$ and therefore $f^*\left(\frac{1}{2} + it\right) \in H_0$.

We note that the RH is equivalent to an order of magnitude $O(x^{\frac{1}{2}+\varepsilon})$ ($\varepsilon > 0$) of $M(x) := \sum_{n \leq x} \mu(n)$, ((ApT) p. 301), i.e., that $M \in H_{\frac{1}{2}+\varepsilon}$. From (LaE6), (MiM), (MiM1), we recall the equivalent Farey series based criterion of $M(x) := \sum_{n \leq x} \mu(n) = \sum_{k=1}^{A(x)} \cos(2\pi r_k)$ with $A(x) := \sum_{n \leq x} \varphi(n)$. From the Sobolev embedding theorem we recall, that $H_{\frac{1}{2}+\varepsilon}$ is a sub-space of C^0 (more precisely the space of Hölder continuous functions with exponent $1/2$) and that the Dirac function δ is an element of the dual space $\delta \in H_{-\frac{1}{2}-\varepsilon}$. The link to „a quick distributional way to the Prime Number Theorem (PNT)“, (ViJ) is given by the distributions $\delta, H[\delta] \in H_{-1/2-\varepsilon}$ where $\pi H[\delta(x)] = \frac{1}{x} = \log' \left(\frac{x}{n} \right)$ and $\psi'(x) = \sum_{n \leq x} \Lambda(n) \delta(x-n)$. We further note, that $H_{\alpha+\varepsilon}$ is compactly embedded into H_α .

We mention the Riemann error function

$$\int_x^\infty \frac{dt}{t(t^2-1) \log t} = \frac{1}{2\pi i} \frac{1}{\log x} \int_{a-i\infty}^{a+i\infty} \frac{d}{ds} \left[\frac{\log \Gamma(1+\frac{s}{2})}{s} \right] x^s ds \cong -\frac{1}{2\pi i} \int_{a-i\infty}^{a+i\infty} \log \Gamma(1+\frac{s}{2}) x^s \frac{ds}{s},$$

derived from the $\Gamma(1+\frac{s}{2})$ term of the entire Zeta function ((EdH) 1.16).

The parts (A1-A3) below are also concerned with the Kummer functions are ${}_1F_1\left(1, \frac{3}{2}; -x\right)$ and ${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; -x\right)$, which are connected by the formula ${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; x\right) = e^x {}_1F_1\left(1, \frac{3}{2}; -x\right)$, (AbM) (7.1.21), (GrI) 9.212:

The Kummer function based representations of the $li(x)$ function is given by

$$li(x) = \int_{-}^x \frac{dt}{\log t} dt = -x {}_1F_1(1; 1, -\log x) = Ei(\log x) = -\int_{-\log x}^{\infty} \frac{e^{-t}}{t} dt = \int_{-\infty}^{\log x} \frac{e^t}{t} dt \sim \frac{1}{\log x}$$

with $Ei(x) = \log x + \gamma + \sum_{k=1}^{\infty} \frac{1}{k!} \frac{x^k}{k}$. The related representation of the $\frac{e^x-1}{x}$ function is given by ((LeN) 9.13.7, 9.8.5),

$$li(x) = x {}_1F_1(1, 1; -\log x) \quad , \quad \frac{e^x-1}{x} = {}_1F_1(1, 2; x).$$

In the following lemma some properties of the Kummer function ${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; x\right)$ are summarised

Lemma:

- i) ${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; x\right) = \sum_{k=0}^{\infty} \frac{1}{k+\frac{1}{2}} \frac{x^k}{k!} = \sum_{k=0}^{\infty} \frac{2}{2k+1} \frac{x^k}{k!} = \frac{1}{2} \int_0^1 \frac{e^{xt}}{\sqrt{t}} dt$, ${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; 0\right) = 1$ (LeN) (9.11.1)
- ii) $x e^{-x^2} {}_1F_1\left(\frac{1}{2}, \frac{3}{2}; x^2\right) = x {}_1F_1\left(1, \frac{3}{2}; -x^2\right) = \int_0^{\infty} e^{-t^2} \sin(2xt) dt$
- iii) ${}_1F_1'\left(\frac{1}{2}, \frac{3}{2}; x\right) = \frac{1}{3} {}_1F_1\left(\frac{3}{2}, \frac{5}{2}; x\right)$
- iv) ${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; x\right) + x {}_1F_1'\left(\frac{1}{2}, \frac{3}{2}; x\right) = e^x$
- v) ${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; x^2\right) = \frac{2}{x} \int_0^x e^{t^2} dt$ (AbM) 7.1.5, (GaW)
- vi) $\int_0^{\infty} x^{s/2} {}_1F_1\left(\frac{1}{2}, \frac{3}{2}; -x\right) \frac{dx}{x} = \frac{\Gamma(\frac{s}{2})}{1-s}$ resp. $\frac{1}{2} \int_0^{\infty} x^{s/2} \left[-x {}_1F_1'\left(\frac{1}{2}, \frac{3}{2}; -x\right)\right] \frac{dx}{x} = \frac{\Gamma(1+\frac{s}{2})}{1-s}$ (GrI) 7.612

Remark: The Hilbert transform of the Gaussian function is the Dawson function given by

$$F(x) := e^{-x^2} \int_0^x e^{t^2} dt = \int_0^{\infty} e^{-t^2} \sin(2xt) dt = x {}_1F_1\left(1, \frac{3}{2}; -x^2\right) = x e^{-x^2} {}_1F_1\left(\frac{1}{2}, \frac{3}{2}; x^2\right).$$

Remark: The functions ${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; -z\right)$ and ${}_1F_1\left(1, \frac{3}{2}; z\right)$ possesses the same zeros (SeA), as the related Whittaker function $z^{-3/4} M_{1,1/2}^{3/4}(z)$, which is the solution of the corresponding self-adjoint (ODE) Whittaker operator ((BuH), §17.1).

Putting

$$g(x) := {}_1F_1\left(\frac{1}{2}, \frac{3}{2}; -x\right) + x {}_1F_1'\left(\frac{1}{2}, \frac{3}{2}; -x\right) = \sum_{k=0}^{\infty} \frac{k+1}{k+1/2} \frac{(-x)^k}{k!} = \frac{1}{2\sqrt{x}} \left[\int_0^x e^{-y} \frac{dy}{\sqrt{y}} - \int_0^x \frac{y}{x} e^{-y} \frac{dy}{\sqrt{y}} \right] = \frac{1}{2} \int_0^x \sqrt{\frac{y}{x}} \left(1 - \frac{y}{x}\right) e^{-y} \frac{dy}{y} \geq 0$$

it follows from lemma 5 v) vii) the

Theorem: In the critical stripe it holds

$$(*) \quad \frac{\Gamma(1+\frac{s}{2})}{s} = \frac{1}{2} \Gamma(\frac{s}{2}) = \frac{1}{s} \int_0^{\infty} x^{s/2} [-x(e^{-x})'] \frac{dx}{x} = \frac{1}{2} \frac{\Gamma(\frac{s}{2})}{1-s} - \frac{\Gamma(1+\frac{s}{2})}{1-s} = \int_0^{\infty} x^{\frac{s}{2}} g(x) \frac{dx}{x}.$$

Therefore, the entire Zeta function $\xi(s) := \frac{s}{2} \Gamma(\frac{s}{2})(s-1)\pi^{-s/2}\zeta(s) = \xi(1-s)$ can be represented as a self-adjoint operator with transform $\xi(s)$ (!!, see also (EdH) 10.3, 10.5)) in the form

$$\xi(s) = s(s-1)\pi^{-s/2}\zeta(s) \left[\int_0^{\infty} x^{\frac{s}{2}} g(x) \frac{dx}{x} \right] = \xi(1-s),$$

whereby both summands of $g(x)$ provide improved asymptotics behavior compared to the Riemann error function (derived from the $\Gamma(1+\frac{s}{2})$ term of the entire Zeta function ((EdH) 1.16)),

$$\int_x^{\infty} \frac{dt}{t(t^2-1)\log t} = \int_x^{\infty} \frac{1}{t\log t} (\sum t^{-2n}) dt = \sum_{n=1}^{\infty} \int_x^{\infty} t^{-2n} \frac{dt}{t\log t} = \frac{1}{2\pi i} \frac{1}{\log x} \int_{a-i\infty}^{a+i\infty} \frac{d}{ds} \left[\frac{\log \Gamma(1+\frac{s}{2})}{s} \right] x^s ds.$$

(*) The new Γ -function representation also enables a new Euler-Mascheroni constant characterization in the form

$$\gamma = \frac{1}{2} \int_0^{\infty} g(x) \log\left(\frac{1}{x}\right) dx.$$

Lemma: (OsH) Vol1, 89): Let $T(x)$ monotone increasing, and

$$f(s) = \int_0^\infty e^{-sx} dT(x) = \int_0^\infty x^{-s} dT(\log x) = \int_0^\infty x^{-s} \frac{dT(x)}{x}, \quad s = \sigma + it, \quad \sigma > 0$$

convergent. Then, if $\lim_{s \rightarrow 0^+} sf(s)$ exists, this holds also for $\lim_{s \rightarrow \infty} \frac{T(x)}{x}$ and both limits are identical, i.e.

$$\lim_{s \rightarrow 0^+} sf(s) = \lim_{s \rightarrow \infty} \frac{T(x)}{x}.$$

Putting

$$T(x) := \int_0^\infty g(t) dt \quad (\text{i.e. } dT(x) = g(x) dx)$$

it follows from the above that $T(x)$ monotone increasing and $\lim_{s \rightarrow \infty} \frac{T(x)}{x} = 1$; for the context to the connection between $\zeta(s)$ and primes resp. Polya's „real self-adjoint operator“ theorem we refer to (EdH) 1.11 resp. (EdH) 12.5, (PoG).

The following part (A1) is related to RH criteria which can be re-formulated in terms of distributional Hilbert scale functions $H_\alpha(-\infty, \infty)$.

Part (A2) is related to RH criteria which can be re-formulated in terms of periodical distributional Hilbert scale functions $H_\alpha^\#(0,1)$.

Part (A3) is about RH criteria formulated by distributional arithmetical functions going beyond the distributional arithmetical functions applied for "a *distributional way to prove the Prime Number Theorem*" (ViJ).

The parts (A2) and (A3) are concerned with appropriate sequences of real numbers and sequences of vectors; the latter ones are called Riesz sequences, if the sequence is a Riesz basis in the closure of the space spanned by those vectors.

With respect to the Bagchi criterion below we emphasize that there is only a positive answer to the following question in a certain „weak“ sense (SeK): „can every frame of complex exponentials $\{e^{i\lambda_n x}\}$ in $L_2(-\pi, \pi)$ be made into a Riesz basis by removing from $\{e^{i\lambda_n x}\}$ a suitable collection of the functions $e^{i\lambda_n x}$ resp. can every Riesz sequence $\{e^{i\lambda_n x}\}$ in $L_2(-\pi, \pi)$ be made into a Riesz basis by adjoining to $\{e^{i\lambda_n x}\}$ a suitable collection of exponentials $e^{i\lambda x}$ not elements of $\{e^{i\lambda_n x}\}$?“ For the critical sequence in our cases (part (A2) and (A3)) this is about $\lambda := \beta_1 := \frac{\omega_1 + \omega_2}{2} - \frac{1}{4}$ to ensure a Snirelmann density of $1/2$ of the concerned sequence $\{\beta_n\}_{n \in \mathbb{N}}$ (see also (ReR)).

Following part (A3), there is a section about a new tool set to prove the binary Goldbach conjecture, taking advantage of (A2-A3), e.g. the sequences based on the imaginary parts of the zeros of the considered Kummer functions with underlying domains with Snirelmann density of $1/2$. The final tool is about a „2-semi-(truly)-circle (even/odd integer) method“ with an underlying „major/minor arcs I/II“ concept alternatively to the Hardy-Littlewood circle method.

(A1) This class is about RH criteria which can be re-formulated in terms of distributional Hilbert scale functions $H_\alpha(-\infty, \infty)$

The Euler product formula connects the Riemann function $\zeta(s)$ (the analytic continuation of the power function representation beyond the halfplane $Re(s) > 1$) with primes. Taking the log of both sides of the Euler product formula results into a Stieltjes integral representation of the $\log \zeta(s)$ function with a (prime) density $dJ(x)$ in the form ((EdH) 1.11)

$$\log \zeta(s) = s \int_0^\infty x^{-s-1} J(x) dx = \int_0^\infty x^{-s} dJ(x)$$

The entire Zeta function

$$\xi(s) := \frac{s}{2} \Gamma\left(\frac{s}{2}\right) (s-1) \pi^{-s/2} \zeta(s) = \xi(1-s)$$

is basically a product of the three functions $\zeta(s)$, $(s-1)$ and $\Gamma(1 + \frac{s}{2})$, whereby $\Gamma(s)$ denotes the Gamma function ((EdH) 1.13).

The method for deriving the formula for $J(x)$ is basically about the calculation of the Fourier inverses of $\log \xi(s)$, $-\log(s-1)$ and $-\log \Gamma(1 + \frac{s}{2})$. The Fourier inverse of the principle term $-\log(s-1)$ becomes the logarithmic integral representation of the $li(x)$ function. The Fourier inverse of the term $-\log \Gamma(1 + \frac{s}{2})$ leads to the famous Riemann approximation error function between the prime density function $J(x)$ and the $li(x)$ function, ((EdH) 1.16),

$$\int_x^\infty \frac{dt}{t(t^2-1)\log t} = \int_x^\infty \frac{1}{t\log t} (\sum t^{-2n}) dt = \sum_{n=1}^\infty \int_x^\infty t^{-2n} \frac{dt}{t\log t} = \frac{1}{2\pi i} \frac{1}{\log x} \int_{a-i\infty}^{a+i\infty} \frac{d}{ds} \left[\frac{\log \Gamma(1 + \frac{s}{2})}{s} \right] x^s ds$$

An appropriate convergence behavior of the Riemann error function is one of several RH criteria. Let $f(x)$ denote the Gaussian function. Then the entire Riemann Zeta function is given by

$$\xi(s) = \frac{s}{2} \Gamma\left(\frac{s}{2}\right) (s-1) \pi^{-s/2} \zeta(s) = (1-s) \cdot \zeta(s) M[-x f'(x)](s) = \xi(1-s).$$

The Hilbert transform of the Gaussian function is the Dawson function given by

$$F(x) := e^{-x^2} \int_0^x e^{t^2} dt = \int_0^\infty e^{-t^2} \sin(2xt) dt = x {}_1F_1\left(1, \frac{3}{2}; -x^2\right) = x e^{-x^2} {}_1F_1\left(\frac{1}{2}, \frac{3}{2}; x^2\right).$$

Replacing the Gaussian function by its Hilbert transform leads to an alternative entire Zeta function definition $\xi^*(s)$ with same zeros in the critical stripe as $\xi(s)$ resulting into a modified Riemann error function with improved approximation property. It is given by

$$\xi^*(s) := \frac{1}{2} (s-1) \pi^{\frac{1-s}{2}} \Gamma\left(\frac{s}{2}\right) \tan\left(\frac{\pi}{2}s\right) \cdot \zeta(s) = \zeta(s) \cdot M\left[\frac{d}{dx} [-x \cdot f_H(x)]\right](s)$$

resp.

$$\zeta^*(s) := \frac{\cot(\frac{\pi}{2}s)}{1-s} \cdot \zeta(s) = \frac{\tan(\frac{\pi}{2}(1-s))}{1-s} \cdot \zeta(s)$$

with same zeros as $\xi(s)$, as it holds $s(1-s)\xi^*(s)\xi^*(1-s) = \pi \xi(s)\xi(1-s)$. It is basically about a replacement of the term $\pi^{\frac{s}{2}}$ of the term $\pi \Gamma\left(1 + \frac{s}{2}\right) = \frac{\pi}{2} s \Gamma\left(\frac{s}{2}\right)$ by the term $\tan\left(\frac{\pi}{2}s\right) = \cot\left(\frac{\pi}{2}(1-s)\right)$. The method for deriving the formula for $J(x)$ is then about the calculation of the Fourier inverses of $\log \xi^*(s)$, $-\log(s-1)$, $-\log \Gamma\left(\frac{s}{2}\right)$ and $-\log\left(\tan\left(\frac{\pi}{2}s\right)\right) = -\log\left(\cot\left(\frac{\pi}{2}(1-s)\right)\right)$.

With the series expansion of $\tan\left(\frac{\pi}{2}s\right)$ resp. $\frac{\pi}{2}s \cdot \cot\left(\frac{\pi}{2}s\right)$ ((GrI) 1.421) it follows

$$\xi^*(s) = \frac{\xi(s)}{\sqrt{\pi}} \sum_{k=1}^\infty \frac{1}{(k-\frac{1}{2})^2 - (\frac{s}{2})^2} \quad \text{resp.} \quad \xi(s) = \frac{\xi^*(s)}{\sqrt{\pi}} \cdot \frac{2}{\pi} \left[1 + \sum_{k=1}^\infty \frac{s^2}{s^2 - (2k)^2} \right].$$

The entire Zeta function $\xi(s) := \frac{s}{2} \Gamma\left(\frac{s}{2}\right) (s-1) \pi^{-s/2} \zeta(s) = \xi(1-s)$ can be also represented as a self-adjoint operator with transform $\xi(s)$ (!! , see also (EdH) 10.3, 10.5)) in the form

$$\xi(s) = s(s-1) \pi^{-s/2} \zeta(s) \left[\int_0^\infty x^{\frac{s}{2}} g(x) \frac{dx}{x} \right] = \xi(1-s)$$

which is one of the RH criteria (CaD).

(A2) This class is about RH criteria which can be re-formulated in terms of periodical distributional Hilbert scale functions $H_{\alpha}^{\#}(0,1)$

Bagchi's "Hilbert space based reformulation of the Nyman-Beurling RH criterion" (BaB) is based on the (periodical) fractional part $L_2^{\#}$ -function

$$\rho(x) = x - [x] = \frac{1}{2} - \sum_1^{\infty} \frac{\sin 2\pi vx}{\pi v}.$$

Its convergent Mellin transform in the critical stripe defines the zeta function $\zeta(s)$ given by ((TiE) II 2.1),

$$\zeta(s) = s \int_0^{\infty} \frac{[x]-x}{x^s} \frac{dx}{x} = -s \int_0^{\infty} x^{-s} \rho(x) \frac{dx}{x} \quad 0 < \operatorname{Re}(s) < 1.$$

The conceptual challenge becomes obvious condering the general Mellin transform property $M[xh'](s) = -sM[h](s)$ and the related (insufficient) Mellin transform zeta function $\zeta(s)$ representation in the form

$$\zeta(s) = s \int_0^{\infty} \frac{[x]-x+\frac{1}{2}}{x^s} \frac{dx}{x} = -s \int_0^{\infty} x^{-s} \left[\rho(x) - \frac{1}{2} \right] \frac{dx}{x} \quad -1 < \operatorname{Re}(s) < 0.$$

The considered Hilbert space in [BaB] is about of all sequences $a = \{a_n | n \in \mathbb{N}\}$ of complex numbers such that

$$\sum_{n=1}^{\infty} \omega_n |a_n|^2 < \infty \quad \text{with} \quad \frac{c_1}{n^2} \leq \omega_n \leq \frac{c_2}{n^2}$$

which is isomorph to the Hilbert space $H_{-1} \cong l_2^{-1}$. For $\gamma := \{1,1,1,1,\dots\}$ it holds

$$\|\gamma\|_{-1}^2 = \sum_1^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$$

i.e. $\gamma \in l_2^1$ resp. $\gamma \in (l_2^{-1/2})^{\perp}$, with $l_2^{-1} = l_2 \otimes l_2^{\perp}$, $l_2^{-1/2} = l_2^{-1/2} \otimes (l_2^{-1/2})^{\perp}$. For the Zeta function on the critical line it also holds $\mathcal{E} \in H_{-1}(-\infty, \infty) = H_{-1} = H_0 \otimes H_0^{\perp}$, and more specifically, $\mathcal{E}, \gamma \in H_{-1/2}^{\perp}$, i.e. the orthogonal projection of \mathcal{E}, γ onto $H_{-1/2}$ is 0.

Theorem (Bagchi-Nyman criterion, (BaB)): Let

$$\gamma_k := \left\{ \rho\left(\frac{m}{k}\right) \mid m = 1, 2, 3, \dots \right\} \quad \text{for } k = 1, 2, 3, \dots$$

and Γ_k be the closed linear span of γ_k . Then the Nyman criterion states that the following statements are equivalent:

- i) The Riemann Hypothesis is true
- ii) $\gamma \in \bar{\Gamma}_k$.

The verification of the Bagchi criterion is enabled by ergodic theory on a Hilbert space, specific properties of the Mellin and Hilbert transforms regarding the basis functions e_v and the fractional function in combination with non-harmonic Fourier series theory (based on complex exponentials forming a Riesz basis for $L_2^{\#}$, ((DuR), (YoR)) and related approximation theory in Hilbert scales.

The concerned $\gamma := \{1,1,1,1,\dots\} \in l_2^{-1}$ can be represented in the form $\gamma = \left\{ \frac{1}{n} \gamma_1 \right\}_{n \in \mathbb{N}} = e_n^{odd} + e_n^{even}$ with $e_{odd} := (1,0,1,0,1,0,1 \dots)$ and $e_{even} := (1,0,1,0,1,0,1 \dots) + (0,1,0,1,0, \dots)$. The analogue split for γ_k with $k \geq 2$ is given by

$$\begin{aligned} \tilde{\gamma}_{2k} &:= \left\{ \frac{1}{n} \gamma_{2k} \right\}_{n \in \mathbb{N}} = \left\{ \frac{e_n^{odd}}{n} \gamma_{2k} \right\}_{n \in \mathbb{N}} + \left\{ \frac{e_n^{even}}{n} \gamma_{2k} \right\}_{n \in \mathbb{N}} \\ \tilde{\gamma}_{2k+1} &:= \left\{ \frac{1}{n} \gamma_{2k+1} \right\}_{n \in \mathbb{N}} = \left\{ \frac{e_n^{odd}}{n} \gamma_{2k+1} \right\}_{n \in \mathbb{N}} + \left\{ \frac{e_n^{even}}{n} \gamma_{2k+1} \right\}_{n \in \mathbb{N}}. \end{aligned}$$

Let $S_n^{odd} := \operatorname{span}\{\tilde{\gamma}_{2k}\}_{k=1,\dots,n+1}$ and $S_n^{even} := \operatorname{span}\{\tilde{\gamma}_{2k+1}\}_{k=1,\dots,n+1}$ denote the n-dimensional subspaces of $H_{-1/2} \cong l_2^{-1/2}$ and $\gamma_{odd}^{(n)} := P_n(e_n^{odd})$ and $\gamma_{even}^{(n)} := P_n(e_n^{even})$ be the orthogonal projections $P_n : H_{-1} \rightarrow S_n^{odd}, S_n^{even}$, of γ onto S_n^{odd}, S_n^{even} . Then the Bagchi criterion is equivalent to

$$(\gamma - \gamma_{odd}^{(n)}, v)_{-1/2} = (\gamma - \gamma_{even}^{(n)}, v)_{-1/2} = 0, \quad \forall v \in l_2^{-1/2}.$$

Lemma ([GrI] 3.761): For $a > 0$ it holds

$$\begin{aligned} M[\sin(a \cdot)](s) &= \int_0^\infty x^s \sin(ax) \frac{dx}{x} = \frac{\Gamma(s)}{a^s} \sin\left(\frac{\pi}{2}s\right) & 0 < \operatorname{Re}(s) < 1 \\ M[\cos(a \cdot)](s) &= \int_0^\infty x^s \cos(ax) \frac{dx}{x} = \frac{\Gamma(s)}{a^s} \cos\left(\frac{\pi}{2}s\right) & 0 < \operatorname{Re}(s) < 1 \end{aligned}$$

and $H[\sin](\omega x) = -\cos(\omega x)$ resp. $H[\cos](\omega x) = \sin(\omega x)$.

The ergodic theorem on a Hilbert space is about the convergent sequence $x_k := \frac{1}{k}(T^1 x + T^2 x + \dots + T^k x)$ with respect to the norm topology. The underlying operator

$$T^* := \lim_{k \rightarrow \infty} \frac{1}{k}(T^1 + T^2 + \dots + T^k)$$

is bounded and fulfills $T^k \circ T^* = T^*$, $T^* \circ T^k = T^*$, $T^* \circ T^* = T^*$. „The mean ergodic theorem ((HaP) p. 16) is an amusing and simple piece of classical analysis in case the underlying Hilbert space is one-dimensional“ (HaP) p. 14.

Lemma ((HiF)): Let E denote the eigen-space of the operator T^* with respect to the eigenvalue 1, i.e. $E := \{x \in H | T^* x = x\}$ then it holds

- i) $E = (T^* - \operatorname{Id})^{-1}(0)$ is closed
- ii) the operator T^* is a projector onto the eigen-space E (because of $T^*|_E = \operatorname{Id}|_E$).

The chosen operator $T^1 := H$ is the Hilbert operator (e.g. (LiI1) 11.1)

$$\underline{H}[u](t) := \frac{1}{i} \int_0^1 \cot(\pi(s-t)) u(s) ds, \quad u \in \bigcap_{\alpha \in \mathbb{R}} H_\alpha.$$

For each $\alpha \in \mathbb{R}$ the extended operators $\underline{H}^{(\alpha)}: H_\alpha \rightarrow H_\alpha$ with domain H_α are bounded Fredholm operators of index zero with $\|\underline{H}^{(\alpha)} u\|_\alpha = \|u\|_\alpha$, with kernel $N(\underline{H}^{(\alpha)}) = \operatorname{span}(1)$ (the set of all constants), and their range is $R(\underline{H}^{(\alpha)}) = \{v \in H_\alpha | (v, 1) = 0\}$, ((LiI1) p.316); and for $e_\nu(x) := e^{2\pi i \nu x}$, $\nu \in \mathbb{Z}$ it holds $\underline{H} e_\nu = \operatorname{sign}(\nu) \cdot e_\nu$ (LiI1).

Putting $T^1 := \underline{H}^{(\alpha)}$ one gets with $T^k e_\nu = \begin{cases} \operatorname{sign}(\nu) e_\nu & \text{for } k \text{ odd} \\ e_\nu & \text{for } k \text{ even} \end{cases}$ for $\nu \neq 0$

$$\sum_{i=1}^{2k-1} T^i e_\nu = \begin{cases} 2k-1 & \text{for } \nu > 0 \\ -1 & \text{for } \nu < 0 \end{cases}, \quad \sum_{i=1}^{2k} T^i e_\nu = \begin{cases} 2k & \text{for } \nu > 0 \\ 0 & \text{for } \nu < 0 \end{cases}.$$

Applying the Hilbert operator to the (periodical) fractional part $L_2^\#$ -function ($\alpha = 0$)

$$\rho(x) = x - [x] = \frac{1}{2} - \sum_1^\infty \frac{\sin 2\pi \nu x}{\pi \nu}$$

leads to (*)

$$\sum_{i=1}^{2k-1} T^i [\rho](x) = \begin{cases} \sum_1^\infty \frac{\cos 2\pi \nu x}{\pi \nu} & \text{for } 2k-1 = 1,5,9,13, \dots \\ \sum_1^\infty \frac{\sin 2\pi \nu x}{\pi \nu} & \text{for } 2k-1 = 3,7,11, \dots \end{cases}$$

and

$$\sum_{i=1}^{2k} T^i [\rho](x) = \begin{cases} \sum_1^\infty \frac{\sin 2\pi \nu x + \cos 2\pi \nu x}{\pi \nu} & \text{for } 2k = 2,6,10 \dots \\ 0 & \text{for } 2k = 4,8,12, \dots \end{cases}$$

whereby $\sum_1^\infty \frac{\cos 2\pi \nu x}{\pi \nu} = -\frac{1}{\pi} \log |2 \sin(\pi x)| =: \rho_H(x) \in L_2^\#(0,1)$ and $\sum_1^\infty \frac{\sin 2\pi \nu x}{\pi \nu} = \frac{1}{2} - \rho(x)$ (**).

We note that the first derivative of the Hilbert transform of $\rho(x)$ is given by the divergent (Cesàro summable) Fourier series representation of the $\cot(\pi x)$ function, $\cot(\pi x) = 2 \sum_1^\infty \sin(2\pi \nu x) = -\rho_H'(x)$ (see also (BeB)); it is an element of the Hilbert space $H_{-1}^\#(0,1)$, i.e. the formulas above are valid in similar form for $\alpha = -1$ applied to the (in $H_{-1}^\#(0,1)$ convergent) Fourier series representation $\frac{1}{2} \cot(\pi x) = \frac{a_0}{2} + \sum_1^\infty a_\nu \cos(2\pi \nu x) + b_\nu \sin(2\pi \nu x)$ with $a_\nu = 0$ for $\nu \geq 0$ and $b_\nu = 1$ for $\nu > 0$ (i.e. $\gamma = \{b_1, b_2, b_3, \dots\}$).

(*) see also (CoG):

$$\begin{aligned} T^1[\rho](x) &= \sum_1^\infty \frac{\cos 2\pi \nu x}{\pi \nu}, \quad T^2[\rho](x) = \sum_1^\infty \frac{\sin 2\pi \nu x}{\pi \nu}, \quad T^3[\rho](x) = -\sum_1^\infty \frac{\cos 2\pi \nu x}{\pi \nu} = -T^1[\rho](x), \quad T^4[\rho](x) = -\sum_1^\infty \frac{\sin 2\pi \nu x}{\pi \nu} = -T^2[\rho](x), \quad T^5[\rho](x) = \\ \sum_1^\infty \frac{\cos 2\pi \nu x}{\pi \nu} &= T^1[\rho](x), \quad T^6[\rho](x) = \sum_1^\infty \frac{\sin 2\pi \nu x}{\pi \nu} = T^2[\rho](x), \quad T^7[\rho](x) = -\sum_1^\infty \frac{\cos 2\pi \nu x}{\pi \nu} = -T^1[\rho](x), \quad T^8[\rho](x) = -\sum_1^\infty \frac{\sin 2\pi \nu x}{\pi \nu} = -T^2[\rho](x), \dots \\ (*[x] + [x + \frac{1}{2}] &= [2x] \text{ resp. more general } \rho(nx) = \sum_{k=0}^{n-1} \rho\left(x + \frac{k}{n}\right), \text{ (ApT) p. 72} \end{aligned}$$

The sequence ω_n (the imaginary parts of the Kummer function ${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; 2\pi iz\right)$) enables the definition of two sequences $\lambda_n^{(1)} = \omega_n$ and $\lambda_n^{(2)} = \frac{\omega_n + \omega_{n+1} - 1}{2}$, leading to „nearly“ harmonic even and odd winding numbers of the corresponding exponentials $\{e^{2\pi i \lambda_n x}\}_{n=0, \pm 1, \pm 2, \dots}$ occurring on the left and right part of the unit circle. From (YoR) p. 36, we recall

Kadec's $\frac{1}{4}$ -Theorem (14): If σ_ν is a sequence of real numbers for which

$$|\sigma_\nu - \nu| \leq L < \frac{1}{4}, \quad \nu = 0, \pm 1, \pm 2, \dots$$

Then $\{e^{i\sigma_\nu x}\}_{\nu=0, \pm 1, \pm 2, \dots}$ satisfies the Paley-Wiener criterion and so forms a Riesz basis for $L_2^\#(-\pi, \pi) = H_0^\#(-\pi, \pi)$.

For the extension of Kadec's theorem to frames (being built by two real sequences $\{\theta_k^{(1)}\}_{k \in \mathbb{Z}}$ $\{\theta_k^{(2)}\}_{k \in \mathbb{Z}}$) we refer to (ChO) 9.8.

If the sequence σ_ν is symmetric, then the product $\prod\left(1 - \frac{z^2}{\sigma_n^2}\right)$ converges to an entire function $f(z)$ which belongs to the Paley-Wiener space, (YoR) p. 124.

The exponentials $e^{i\sigma_\nu x}$ can be transformed into the reproducing functions

$$K_n(z) = \frac{\sin(\pi(z - \sigma_n))}{\pi(z - \sigma_n)}.$$

For $G(z) := z \prod\left(1 - \frac{z^2}{\sigma_n^2}\right)$ the related function $G_n(z) := \frac{G(z)}{G'(\sigma_n)(z - \sigma_n)}$ can be transformed into $g_n(t) := \int_{-\infty}^{\infty} G_n(x) e^{ixt} dt$. The solution of the moment problem $f(\sigma_\nu) = c_\nu$ $\nu = 0, \pm 1, \pm 2, \dots$, is given by (YoR) p.126,

$$(*) \quad f(z) = \sum_{\nu=-\infty}^{\infty} c_\nu \frac{G(z)}{G'(\sigma_\nu)(z - \sigma_\nu)}.$$

Remark (YoR) p.126: The formula (*) is a simple example of a „generalized“ Lagrange interpolation formula for an entire function assuming the values c_ν at the points σ_ν .

Remark: The product representation of the considered Kummer function is given by ((BuH) p.184)

$${}_1F_1(a, c, z) = \frac{1}{\Gamma(c)} e^{\frac{a}{c}z} \prod\left(1 - \frac{z}{\alpha_n}\right) e^{z/\alpha_n}.$$

The considered imaginary parts of its zeros fulfill the Kadec requirements with

$$\sigma_n = \alpha_n := \omega_n + \frac{1}{4}, \quad \sigma_n = \beta_n := \frac{\omega_n + \omega_{n+1}}{2} - \frac{1}{4}.$$

We mention that the approximation solution to $\cot(\pi x)$, being testing against $H_0^\#(0,1)$ test functions, is an element of $H_{-1/2}^\#(0,1)$.

With respect to (A1) and the newly proposed entire Zeta function $\xi^*(s) := \frac{1}{2}(s-1)\pi^{\frac{1-s}{2}}\Gamma\left(\frac{s}{2}\right)\tan\left(\frac{\pi}{2}s\right) \cdot \zeta(s)$ we note the related (classical) series representations for the $\log\left(\tan\left(\frac{\pi}{2}x\right)\right)$ (whereby all summands of the first series are positive), and the (A2)-related series representations for the $\log \sin(\pi x)$ functions, given by (GrI) 1.518,

$$\log\left(\tan\left(\frac{\pi}{2}x\right)\right) = \log\left(\frac{\pi}{2}x\right) + \sum_{k=1}^{\infty} (-1)^{k+1} \frac{1}{k} \frac{(2^{2k-1}-1)}{(2k)!} B_{2k}(\pi x)^{2k}, \quad x^2 < 1$$

$$\log \sin(\pi x) = \log(x\pi) + \sum_{k=1}^{\infty} (-1)^k \frac{1}{k} \frac{2^{2k-1}}{(2k)!} B_{2k}(\pi x)^{2k}, \quad x^2 < 1.$$

The term $-\log\left(\tan\left(\frac{\pi}{2}s\right)\right)$ is $L_2^\#$ -integrable and can be represented as a convergent series (EIL). It results into a correspondingly modified Riemann error function or a correspondingly modified definition of the $li(x)$ function. In both cases the approximation behavior between the $li(x)$ function and the prime density function gets improved.

(A3) This class is about RH criteria formulated by distributional arithmetical functions going beyond the distributional arithmetical functions applied for "a distributional way to prove the Prime Number Theorem" (ViJ).

In 1927 Polya published a very different sort of theorem as in (PoG) on the same general subject ("Über trigonometrische Integrale mit nur reellen Nullstellen") with quite weak conditions to the baseline function $F(u)$ (see also (EdH) 12.5). All attempts failed so far to build a proper (classical) baseline function $F(u)$ to prove the RH. The proposed distributional Hilbert space framework of this homepage provides an additional opportunity to build such a baseline function.

According to E. Landau the classical proofs of the Prime Number Theorem (PNT) are not being scalable for "deeper going number theory problems". Ikehara's theorem ((EdH) 12.7) is about a deduction from Wiener's general Tauberian theorem. In (ViJ) "a distributional way to prove the PNT" is based on the Mellin transform of e.g. the Dirac delta "function". The part B section is about a (distributional) quantum element/energy Hilbert space framework enabling a quantum gravity model. From a physical modelling perspective it goes along with a replacement of Dirac's model of the "density" of an "idealized point mass" resp. an "idealized point charge" (modelled by the Dirac or Delta "function") by Plemelj's concept of a "mass element dm " (PIJ) of the proposed "bosons" quantum state Hilbert space $H_{-1/2}$. "Tauberian Theorems for Generalized Functions" come along with the concept of an "automodel function". It is claimed to be an appropriate framework to apply the Polya theorem to prove the RH, as the crucial condition of Polya's theorem is equivalent to one of the characterization criteria of an "automodel function", ((VIV) p. 57).

The proposed periodical, distributional Hilbert scale framework $H_\alpha^\#$ also enables also a truly unit circle method to tackle the binary Goldbach problem. It overcomes current handicaps of the famous Hardy-Littlewood circle method, which is based on the open unit circle disk.

Vinogradov applied the Hardy-Littlewood circle method to derive his famous (currently best known, but not sufficient) estimate regarding the tertiary Goldbach problem. Vinogradov's theorem states that any sufficiently large odd integer can be written as a sum of three prime numbers. The "any sufficiently large odd integer" condition is due to a not sufficiently "good" estimate: it is derived from two components based on a decomposition of the (Hardy-Littlewood) circle into two parts, the "major arcs" (also called "basic intervals") and the "minor arcs" (also called "supplementary intervals"). The sufficiently good estimate is based on "major arcs" estimate using also Goldbach problem relevant data; the not sufficiently good "minor arcs" estimate are purely Weyl sums estimates taking not any Goldbach problem relevant data into account. However, this estimate is optimal with respect to Weyl sums properties. In other words, the major/minor arcs decomposition is inappropriate to solve both Goldbach problems. The proposed periodical, distributional Hilbert scale framework $H_\alpha^\#$ with its underlying domain (the boundary of the unit disk, the unit circle) is claimed to enable a truly unit circle based method to prove the binary Goldbach problem.

The proposed change also enables the definition of *two* appropriately different arithmetical functions to analyze binary number theoretical problems for any prime number pair (p, q) . We note that the winding numbers (i.e. the set of integers) of the unit circle are related to the zeros of the Weyl sum components, which are the basis functions $e^{2\pi i n x}$. The sequence ω_n (the imaginary parts of the Kummer function ${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; 2\pi i z\right)$) enables the definition of two sequences $\lambda_n^{(1)} = \omega_n$ and $\lambda_n^{(2)} = \frac{1}{2}(\omega_n + \omega_{n+1})$, leading to "nearly" harmonic even and odd winding numbers of the corresponding exponentials $\{e^{2\pi i \lambda_n x}\}_{n=0, \pm 1, \pm 2, \dots}$ occurring on the left and right semicircle of the unit circle.

Referring back to (A2) and the considered $\gamma = \{1, 1, 1, 1, \dots\}$ with $\gamma \in (L_2^{-1/2})^\perp$ we mention that γ can be interpreted as a "winding number list", while running periodically through the unit circle

The links of non-harmonic Fourier series theory to the cardinal series based interpolation theory in the Paley-Wiener space PW are given by the following theorems (YoR) p. 170, p. 157):

for a complex exponentials system $\{e^{2\pi i \lambda_n x}\}_{n=0, \pm 1, \pm 2, \dots}$ to form a Riesz basis for $L_2^\#$ it is necessary and sufficient to the interpolation problem $f(\lambda_n) = c_n$, where $f \in PW$

a sequence of vectors φ_n belonging to a separable Hilbert space H is a Riesz basis if and only if it is an exact frame, i.e. $A\|f\|^2 \leq \sum_{n=1}^{\infty} |(f, \varphi_n)|^2 \leq B\|f\|^2$ (if φ_n is a complete orthogonal sequence in H , then it holds $A = B = 1$), or if and only if φ_n is both, a frame and a Riesz sequence (SeK)

The link to a specific class of non-harmonic Fourier series is given by the considered sequences of real or complex numbers with uniform density one (DuR).

A tool set to prove the binary Goldbach conjecture

On the occasion of the 59th birthday
of my wife, Vibhuta
August 25, 2020

In the following we summaries the proposed tool set to tackle the binary Goldbach conjecture.

The following two lemmata are about the building of arithmetical functions based on a certain integrals:

Lemma 1 (Landau, (PoG1)): Let q_n denote a divergent sequence of positive numbers $0 < q_1 \leq q_2 \leq q_3 \leq \dots$ $\lim_{n \rightarrow \infty} q_n = \infty$, $\tau(x)$ the corresponding counting function of the numbers of q_n less than $\leq x$ and $w(x)$ a positive, non-decreasing function with $\lim_{x \rightarrow \infty} \frac{w(2x)}{w(x)} = \lim_{x \rightarrow \infty} \frac{\tau(x)w(x)}{x} = 1$.

Then

$$\lim_{x \rightarrow \infty} \frac{1}{\tau(x)} \sum_{q \leq x} \rho\left(\frac{x}{q}\right) = 1 - \gamma,$$

where $\rho(x)$ denotes the fractional part function.

In (PoG1) lemma 1 is generalized by

Lemma 2: Let $w(x)$ a positive, non-decreasing function with $\lim_{x \rightarrow \infty} \frac{w(\beta x)}{w(\alpha x)} = 1$ with α, β positive numbers. Then

$$\lim_{x \rightarrow \infty} \frac{w(x)}{x} \sum_{n \leq x} f\left(\frac{x}{n}\right) = \int_0^1 f(t) dt.$$

Remark 1: Lemma 2 is valid for two kind of conditions about $f(t)$:

- i) $f(t)$ is Riemann integrable in $[0,1]$
- ii) $f(t)$ is integrable in every closed sub-interval of $[0,1]$, which does not contain 0, and it exists an $\alpha \in (0,1)$ with $\lim_{t \rightarrow 0} t^{1-\alpha} f(t) = 0$.

Remark 2: The condition $\lim_{x \rightarrow \infty} \frac{w(\beta x)}{w(\alpha x)} = 1$ is related to the theory of quasi-asymptotics of generalized functions with its underlying concept of (slow) regular varying (automodel) functions $\vartheta(x)$, where $\lim_{x \rightarrow \infty} \frac{\vartheta(\beta x)}{\vartheta(x)}$ (resp. $\lim_{x \rightarrow \infty} \frac{x\vartheta'(x)}{\vartheta(x)}$) exists, (VIV) p. 56/57; (EsR). The latter condition, $\lim_{x \rightarrow \infty} \frac{x\vartheta'(x)}{\vartheta(x)} = c$, links back to the famous Polya criterion in (PoG) given by $\alpha \leq -\frac{xf'(x)}{f(x)} \leq \beta$ being valid for only (!) closed interval domains. We note that $-\log x$ is slowly varying at $x = 0^+$, (SeE) p.47. We further note with respect to lemma 5 v) below, that for $\tilde{h}(x) := {}_1F_1\left(\frac{1}{2}, \frac{3}{2}; x\right)$ it holds $-\frac{x\tilde{h}'(x)}{\tilde{h}(x)} = 1 - \frac{e^x}{h\tilde{h}x}$.

In (LaE) §56, the following following lemma is provided. It has been applied in (LaE1) to derive the asymptotics integral formula for the Goldbach number counting function

$$H(2n) := \sum_{k=1}^{2n} G_{2k} = \sum_{p \leq 2n} \pi(2n - p), \quad p + q \leq 2n,$$

in the form

$$H(2n) \sim 2n \int_2^n \frac{1}{\log(2n-u) \log u} du \sim 2n \int_2^n \frac{1}{\log(2n-\frac{u}{2}) \log u} du \sim \frac{1}{2} \left(\frac{2n}{\log(2n)}\right)^2, \quad 0 \leq \vartheta \leq 1$$

by putting $F(u, x) := \pi(2n - p)$.

We note that $\sum_{k \text{ odd}}^{2x} G_k \sim \frac{2x}{\log(2x)}$, while $\sum_{k=1}^x G_{2k} \sim \frac{1}{2} \left(\frac{2x}{\log(2x)}\right)^2$, because the odd k can be neglected as every odd number can be represented as sum of two primes, if $k - 2$ is prime, otherwise not (LaE1).

We further note that $d\left[\sum_{k=1}^x \frac{1}{\varphi(k)}\right] \sim d[\log x] \sim \frac{dx}{x}$ (LaE1), and $d\vartheta = \left[\sum_{n \leq x} \frac{\mu(n)}{n} \log \left(\frac{x}{n}\right)\right] = \frac{1}{x} \left[\sum_{n \leq x} \frac{\mu(n)}{n}\right] dx \sim \frac{dx}{x}$ (ApT) p.97.

Lemma 3: Let $F(u, x)$ be a function with real arguments with $2 \leq u \leq x$ fulfilling the following conditions

- i) $F(u, x) \geq 0$
- ii) For fixed $x > 2$ the function $\frac{F(u, x)}{\log u}$ is never increasing for $u \in (2, x)$
- iii) $F(2, x) = o\left(\int_2^x \frac{F(u, x)}{\log u} du\right)$,

then it holds

$$\sum_{p \leq x} F(p, x) \sim \int_2^x \frac{F(u, x)}{\log u} du .$$

The considered Stäckel approximation formula in (LaE1) is given by

$$\tilde{H}(2n) := \sum_{k=1}^n \tilde{G}_{2k} = c_\lambda \cdot \sum_{\substack{k=2 \\ k \text{ even}}}^{2n} \frac{1}{\varphi(k)} \left(\frac{k}{\log(k)}\right)^2 \quad \text{with } c_\lambda := \frac{\pi^4}{105 \cdot \zeta(3)} \sim 0,772 \dots$$

The factor c_λ was suggested to ensure that $\tilde{H}(2n) < H(2n)$.

Remark 3: We note that the above (appreciated) identical asymptotical behavior of both summation formulas does not allow a corresponding conclusion to the approximation behavior of the underlying Goldbach numbers G_{2k} resp. \tilde{G}_{2k} , as in the sum formula $\sum_{k=1}^n G_{2k} - \tilde{G}_{2k}$ positive and negative terms cancel each other out.

Remark 4: With respect to (A2) above and the the term $\frac{1}{\varphi(n)}$ in the Stäckel approximation formula we note the inequality, (ApM) p. 71, $\frac{\sigma(k)}{k^2} \leq \frac{1}{\varphi(k)} \leq \frac{\pi^2}{6} \frac{\sigma(k)}{k^2} = \zeta(2) \frac{\sigma(k)}{k^2}$, $k \geq 2$, where $\sigma(n) = \sigma_1(n)$ denotes the sum of the divisors of n ((ApM) p. 38), and $\varphi(n)$ denotes the Euler totient function. It can be represented as discrete Fourier transform of the greatest common divisor in the form $\varphi(n) = \sum_{k=1}^n \gcd(k, n) \cdot \cos(2\pi \frac{k}{n})$. We further mention that for even integers it holds $\varphi(2k) = 2\varphi(k) > \sqrt{k}$. At the same point in time, it holds $\frac{k}{\log^2 k} > \sqrt{k}$ for $k \geq 2$.

Remark 5: It holds $\sum_{k \leq x} \frac{1}{\varphi(k)} = O(\log x)$, $\frac{1}{x} \sum_{k \leq x} d(n) = \log x + (2\gamma - 1) + O(\sqrt{x})$, and $\sum_{k \leq x} \frac{k}{\varphi(k)} = O(x)$, $\sum_{k \leq x} \frac{1}{\varphi(k)} = O(\log x)$ (ApM) p. 71. The average order of $d(n)$ is $\log n$ i.e. $\frac{1}{x} \sum_{k \leq x} d(k) = \log x + 2\gamma - 1 + O(\sqrt{x}) \sim \log x$, (ApM) p. 52. For $\vartheta(x) := \sum_{n \leq x} \frac{\mu(n)}{n} \log \left(\frac{x}{n}\right)$ we recall from (ApT) p. 97, that the PNT theorem is equivalent to $d\vartheta(x) = \sum_{n \leq x} \frac{\mu(n)}{n} \frac{dx}{x} \sim \frac{dx}{x}$.

Remark 6: ((LaE) §56 p. 214): the n -th prime number is asymptotically equal to $n \cdot \log n$.

Building an arithmetical function for the term $F(u, x) := \pi(2n - p)$ is challenging because of

Lemma 4, ((LaE) §56, p. 215):

from a certain number x on, there are more primes in the interval $(1, x)$ than in the interval $(x, 2x)$.

Remark 7: Lemma 4 cannot be proven with the PNT; the proof requires the asymptotics

$$\pi(x) = \frac{x}{\log x} + \frac{x}{\log^2 x} + o\left(\frac{x}{\log^2 x}\right)$$

Remark 8: The functions ${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; -z\right)$ and ${}_1F_1\left(1, \frac{3}{2}; z\right)$ possesses the same zeros as the related Whittaker function $z^{-3/4} M_{\frac{1}{4}, \frac{1}{4}}(z)$, which is the solution of the corresponding self-adjoint (ODE) Whittaker operator ((BuH), §17.1). The latter one also plays key role for the radial Schrödinger operator with a Coulomb potential or with a morse potential on the half line (DeJ), (LaJ1).

Corresponding integrals of oscillatory type for ${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; x\right)$ (when the variables are real) are the Fresnel integrals $C(x), S(x)$ with $\lim_{x \rightarrow \infty} C(x) = \lim_{x \rightarrow \infty} S(x) = \frac{1}{2}$ with the exact error formula (OIF) p. 67,

$$C\left(\sqrt{\frac{2x}{\pi}}\right) + i \cdot C\left(\sqrt{\frac{2x}{\pi}}\right) = \sqrt{\frac{2x}{\pi}} {}_1F_1\left(\frac{1}{2}, \frac{3}{2}; ix\right) = \int_0^x \frac{\cos t dt}{\sqrt{2\pi t}} + i \cdot \int_0^x \frac{\sin t dt}{\sqrt{2\pi t}} = \left(\frac{1}{2} + i\frac{1}{2}\right) - \frac{ie^{ix}}{\sqrt{2\pi x}} \sum_{n=0}^{\infty} \frac{1 \cdot 3 \cdot \dots \cdot (2n-1)}{(2ix)^n}.$$

The formula allows an alternative „2-semicircle (even/odd integer) method“ with an underlying „major/minor arcs I/II“ concept based on two appropriately defined Fresnel integral distribution functions ($\cos(t) > 0, \sin(t) > 0$), enabled by the real part values ω_n of ${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; 2\pi iz\right)$. With respect to the link to exponentials Riesz bases we refer to (KoG), (NaA) and the references cited there.

Lemma 6:

For the real part values ω_n of ${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; 2\pi iz\right)$ it holds (SeA)

- i) $2n - 1 < 2\omega_n, \omega_n + \omega_{n+1} - 1 < 2n < 2\omega_n + 1, \omega_n + \omega_{n+1} < 2n + 1$
- ii) $n - \frac{3}{4} < \lambda_n^{(1)} - \frac{1}{4} := \omega_n - \frac{1}{4}, \lambda_n^{(2)} - \frac{1}{4} := \frac{\omega_n + \omega_{n+1} - 1}{2} - \frac{1}{4} < n - \frac{1}{4} < \omega_n + \frac{1}{4}, \frac{\omega_n + \omega_{n+1} - 1}{2} + \frac{1}{4} < n + \frac{1}{4}$
- iii) $n - \frac{1}{2} < \omega_n < n, \frac{1}{2} < \omega_1, \frac{\omega_1 + \omega_2 - 1}{2} < 1, s_n := \frac{\omega_n}{n} \rightarrow 1 \quad n \in \mathbb{N}$
- iv) the non-integer sequences $2\omega_n$ and $\omega_n + \omega_{n+1}$ fulfill a kind of Hadamard gap condition

$$\frac{\omega_{n+1}}{\omega_n} > \frac{n + \frac{1}{2}}{n} = 1 + \frac{1}{2n} > q > 1 \quad \text{resp.} \quad \frac{\omega_{n+1} + \omega_{n+2}}{\omega_n + \omega_{n+1}} > \frac{2n+2}{2n+1} = 1 + \frac{1}{2n+1} > q > 1$$
- v) $\theta := \frac{1}{4} < \omega_{n+1} - \omega_n < 1 - \frac{1}{4} = 1 - \theta$
- vi) putting $a_n := \frac{\omega_n}{n} - \frac{1}{2}, b_n := \frac{\omega_1 + \omega_2 - 1}{2n} - \frac{1}{2}$ it follows

$$0 < a_1 = \omega_1 - \frac{1}{2} \leq a_n \rightarrow \frac{1}{2}, \quad \frac{1}{2} \leftarrow b_n < b_1 = \frac{\omega_1 + \omega_2 - 1}{2} < 1, \quad a_n, b_n - a_n \in \left(0, \frac{1}{2}\right), \quad b_n, 1 - a_n \in \left(\frac{1}{2}, 1\right).$$

Lemma 7, (BaR), (KaD):

The following inequalities are valid

$$c_1 \cdot \left[{}_1F_1\left(\frac{1}{2}, c; x\right) \right]^2 \leq {}_1F_1(a_n, c; x) \cdot {}_1F_1(1 - a_n, c; x) \leq \left[{}_1F_1\left(\frac{1}{2}, c; x\right) \right]^2.$$

Remark 9: For the sequences $a_n := \frac{\omega_n}{n} - \frac{1}{2}, b_n := \frac{\omega_n + \omega_{n+1} - 1}{2n} - \frac{1}{2}$ it holds $a_n \in \left(0, \frac{1}{2}\right)$ resp. $b_n \in \left(\frac{1}{2}, 1\right)$ with $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = 1/2$, and the domain of each sequence has Snirelmann density $\frac{1}{2}$. Therefore, it holds

$$-x {}_1F_1(a_n; a_n + 1, -\log x), -x {}_1F_1(b_n; b_n + 1, -\log x) \rightarrow li(x) = -x {}_1F_1(1; 1, -\log x).$$

Remark 10: In the context of non-harmonic Fourier series theory the counterpart of $\gamma := \{1, 1, 1, \dots\}$ can be defined by $(a, b)_n := \{a_1, b_1, a_2, b_2, \dots\}$. It enables „two check points“, while running once through the unit circle, one „check point“ for each prime number p, q .

Remark 11: The sequences $s_n := \frac{\omega_n}{n}$ and $t_n := \frac{\omega_n + \omega_{n+1} - 1}{2n}$ fulfill the Hardy-Littlewood condition $|s_{n+1} - s_n| < \frac{K}{n}$, e.g. s_n has a defined Abel average ((EdH) 12.7) $\lim_{r \uparrow 1} \frac{s_1 r + s_1 r^2 + s_3 r^3 + \dots}{r + r^2 + r^3 + \dots} = L$.

Remark 12 ((KaM), (KoA), (ZyA)): Let $\{n_k\}$ be a sequence of integers satisfying the Hadamard gap condition, i.e. $\frac{n_{k+1}}{n_k} > q > 1$, then trigonometric gap series $\sum_{k=1}^{\infty} c_k \sin(2\pi n_k x)$ converges almost everywhere iff $\sum_{k=1}^{\infty} c_k^2 < \infty$.

Remark 13: We note that for $\tilde{T}(\log x) := \sum_{n \leq x} \frac{1}{n} \log\left(\frac{x}{n}\right), x \geq 1$, the inverse mapping is given by ([ScW] lemma 3.3)

$$\sigma(x) := \tilde{T}^{-1}(\log x) = \sum_{n \leq x} \frac{\mu(n)}{n} \log\left(\frac{x}{n}\right).$$

Remark 14: With respect to Remark 7 we note that what also cannot be derived from the PNT is the convergence of the series

$$\sum_{n=1}^{\infty} \frac{\mu(n)}{n} \log\left(\frac{1}{n}\right) = 1 .$$

„The corresponding theorem goes deeper than the PNT, and from it the PNT can be easily derived“ ((LaE) §160). From (ApT) p. 66 we recall that the PNT can be derived from $\sum_{n=1}^{\infty} \frac{\mu(n)}{n} = 0$.

The related point measure is given by $\vartheta(x) = \sum_{n \leq x} \frac{\mu(n)}{n} \log\left(\frac{x}{n}\right)$ with $d\vartheta = \frac{1}{x} \left[\sum_{n \leq x} \frac{\mu(n)}{n} \right] dx$. Because of $\frac{d}{dx} \log\left(\frac{x}{a}\right) = \frac{1}{x}$ it can be replaced by

$$\vartheta^*(x) = \sum_{\substack{n \leq x \\ n \text{ odd}}} \frac{\mu(n)}{n} \log\left(\frac{2x}{2\omega_n - \frac{1}{2}}\right) + \sum_{\substack{n \leq x \\ n \text{ even} \\ n=1}} \frac{\mu(n)}{n} \log\left(\frac{2x}{\omega_n + \omega_{n+1} - \frac{1}{2}}\right)$$

with $d\vartheta = d\vartheta^* \sim \frac{dx}{x} = d(\log x)$ resp.

$$\sum_{\substack{n \leq x \\ n \text{ odd}}} \frac{\mu(n)}{n} \log\left(\frac{2}{2\omega_n - \frac{1}{2}}\right) + \sum_{\substack{n \leq x \\ n \text{ even} \\ n=1}} \frac{\mu(n)}{n} \log\left(\frac{2}{\omega_n + \omega_{n+1} - \frac{1}{2}}\right) \leq 1$$

The additional term $\omega_n + \omega_{n+1} - \frac{1}{2}$ for the „ n even“ series (i.e. not starting the series $\omega_n + \omega_{n+1} + \frac{1}{2}$) is to get „ $n = 1$ “ as an element of the underlying domain, i.e. both domains get a Snirelmann density of $\frac{1}{2}$. Additionally both related sequences, $\left(\frac{2}{2\omega_n - \frac{1}{2}}\right)^{-1} = \omega_n - \frac{1}{4}$ and $\left(\frac{2}{\omega_n + \omega_{n+1} - \frac{1}{2}}\right)^{-1} = \frac{\omega_n + \omega_{n+1}}{2} - \frac{1}{4}$ fulfill the prerequisite of the ((YoR) p. 36)

Kadec $\frac{1}{4}$ -Theorem: If σ_ν is a sequence of real numbers for which

$$|\sigma_\nu - \nu| \leq L < \frac{1}{4}, \quad \nu = 0, \pm 1, \pm 2, \dots,$$

then $\{e^{i\sigma_\nu x}\}_{\nu=0, \pm 1, \pm 2, \dots}$ satisfies the Paley-Wiener criterion and so forms a Riesz basis for $L^2(-\pi, \pi) = H_0^\#(-\pi, \pi)$.

The below strongly increasing sequences $(A_n^{(1)}, \bar{A}_n^{(1)})$ and $(A_n^{(2)}, \bar{A}_n^{(2)})$ (lemma 9) enable the definition of a twofold finer sieve method (HaH): there is the split above between „odd“ and „even“ series, both with underlying Snirelmann density $\frac{1}{2}$, governing two different arithmetical functions on the first level; on a second level per each series there can be built specific sieve methods per underlying considered „prime interval“ domains $[1, p]$ and $[p, 2p]$, to overcome the mentioned challenge in lemma 4 to build an arithmetical function for $F(u, x) := \pi(2n - p)$ based on lemma 2.

From (LaE1) we note that $\sum_{k \text{ odd}}^{2x} G_k \sim \frac{2x}{\log(2x)}$, while $\sum_{k=1}^x G_{2k} \sim \frac{1}{2} \left(\frac{2x}{\log(2x)}\right)^2$, because the odd k can be neglected as every odd number can be represented as sum of two primes, if $k - 2$ is prime, otherwise not.

Lemma 8:

For the sequences pair $(\lambda_k^{(1)}, \lambda_k^{(2)})$ it holds $\lambda_k^{(1)} \in \left(k - \frac{1}{2}, k\right)$, $\lambda_k^{(2)} \in \left(k, k + \frac{1}{2}\right)$, and therefore,

$$\frac{2k-1}{2k+1} < \frac{\lambda_k^{(1)}}{k} \cdot \frac{\lambda_k^{(2)}}{k+\frac{1}{2}} < 1 \quad \text{resp.} \quad \frac{1}{2k} \cdot \frac{1}{2(k+1)} < \frac{1}{2k} \cdot \frac{2k-1}{2k+1} < \frac{1}{2k} \cdot \frac{\lambda_k^{(1)}}{k} \cdot \frac{\lambda_k^{(2)}}{k+\frac{1}{2}} < \frac{1}{2k}$$

resp.

$$\frac{n}{2}n < \sum_{k=1}^n \lambda_k^{(1)} < \frac{n}{2}(n+1) < \sum_{k=1}^n \lambda_k^{(2)} < \frac{n}{2}(n+2).$$

Lemma 9:

Putting $a_k^{(1)} := \lambda_k^{(1)} - \frac{1}{2}$, $a_k^{(2)} := \lambda_k^{(2)} - \frac{1}{2}$ one gets

$$(2n-1) < A_n^{(1)} := \frac{1}{n} \sum_{k=1}^{2n} a_k^{(1)}, \quad \bar{A}_n^{(1)} := \frac{1}{n} \sum_{k=1}^{2n} \left[2n - a_k^{(2)}\right] < 2n < \\ < A_n^{(2)} := \frac{1}{n} \sum_{k=1}^{2n} a_k^{(2)}, \quad \bar{A}_n^{(2)} := \frac{1}{n} \sum_{k=1}^{2n} \left[2n - a_k^{(1)}\right] < (2n+1)$$

and each of the two domains of the two merged, strongly increasing sequences $(A_n^{(1)}, \bar{A}_n^{(1)})$ and $(A_n^{(2)}, \bar{A}_n^{(2)})$ both have Snirelmann density $\frac{1}{2}$.

The link to the proposed Kummer function based Zeta function theory is given by the theory of quasi-asymptotics in distributional Hilbert scales, defined via the eigenvalues of the specific Whittaker operator with the fundamental solutions, (AbM) 13.1.31, (GrI) 9.212,

$${}_1F_1\left(\frac{1}{2}, \frac{3}{2}; z\right) = z^{-\frac{3}{4}} e^{\frac{z}{4}} M_{\frac{1}{4}}(z) \quad \text{and} \quad {}_1F_1\left(1, \frac{3}{2}; -z\right) = z^{-\frac{3}{4}} e^{-\frac{z}{4}} M_{\frac{1}{4}}(z).$$

From lemma 5 xi) we recall the asymptotics

$$d\left[{}_1F_1\left(\frac{1}{2}, \frac{3}{2}; \log x\right)\right]^2 \sim \left(\frac{x}{\log x}\right)^2 dx.$$

For the construction of a Kummer function related arithmetical function one gets from lemma 5 vii)) with $s = 1/2$

$$\int_0^\infty t^{1/4} {}_1F_1\left(\frac{1}{2}, \frac{3}{2}; -t\right) \frac{dt}{t} = 2\Gamma\left(\frac{1}{4}\right),$$

i.e. ($t \rightarrow -\log x = \log \frac{1}{x}$) it holds

$$\int_0^1 h(x) dx = 1 \quad \text{for} \quad h(x) := \frac{1}{x} \frac{\log^{\frac{1}{4}}\left(\frac{1}{x}\right)}{2\Gamma\left(\frac{1}{4}\right)} \frac{{}_1F_1\left(\frac{1}{2}, \frac{3}{2}; \log x\right)}{\log x} = \frac{1}{2\Gamma\left(\frac{1}{4}\right)} \frac{1}{\sqrt{x \log x}} \frac{M_{\frac{1}{4}}(\log x)}{\log x}.$$

With the asymptotics, (AbM) 13.1.4, 13.1.5,

$$\text{i)} \quad {}_1F_1'(a, c; z) = \frac{\Gamma(c)}{\Gamma(a)} e^z z^{a-c} [1 + O(|z|^{-1})] \quad , \quad \text{Re}(z) > 0$$

$$\text{ii)} \quad {}_1F_1'(a, c; z) = \frac{\Gamma(c)}{\Gamma(c-a)} (-z)^{-a} [1 + O(|z|^{-1})] \quad , \quad \text{Re}(z) < 0$$

it follows

$$\int_0^1 \frac{d}{dt} \left[{}_1F_1\left(\frac{1}{2}, \frac{3}{2}; \log t\right) \right]^2 dt = \left[{}_1F_1\left(\frac{1}{2}, \frac{3}{2}; 1\right) \right]^2 - \left[{}_1F_1\left(\frac{1}{2}, \frac{3}{2}; -\infty\right) \right]^2 = 1.$$

Remark: Recalling ${}_1F_1(a, a+1; x) \sim \frac{1}{\Gamma(a)} \frac{e^x}{x}$ and $\frac{1}{2} \frac{d}{dx} F_a^2(x) = \frac{a}{a+1} F_a(x) \cdot F_{a+1}(x)$ two (strongly increasing resp. strongly decreasing) sequences $\vartheta_n^{(i)}$ with $0 < \vartheta_n < 1$ and $\lim_{n \rightarrow \infty} \vartheta_n^{(i)} = \frac{1}{2}$ ($i = 1, 2$) allow to approximate the $d\left[{}_1F_1\left(\frac{1}{2}, \frac{3}{2}; \log t\right)\right]^2$ integral density in the following form

$$\begin{aligned} \lim_{n \rightarrow \infty} \left[{}_1F_1'\left(\vartheta_n^{(1)}, \vartheta_n^{(1)} + 1; \log x\right) \cdot {}_1F_1'\left(\vartheta_n^{(2)} + 1, \vartheta_n^{(2)} + 2; \log x\right) \right] &= \frac{1}{2} \frac{d}{dx} \left[{}_1F_1\left(\frac{1}{2}, \frac{3}{2}; \log x\right) \right]^2 \\ \frac{\pi}{2} {}_1F_1'\left(\vartheta_n^{(1)}, \vartheta_n^{(1)} + 1; \log x\right) \cdot {}_1F_1'\left(\vartheta_n^{(2)} + 1, \vartheta_n^{(2)} + 2; \log x\right) &\sim \frac{\pi}{2} \frac{1}{\Gamma(\vartheta_n^{(1)})} \frac{1}{\Gamma(\vartheta_n^{(2)})} \left[\frac{x}{\log x} \right]^2 \sim \frac{1}{2} \left[\frac{x}{\log x} \right]^2. \end{aligned}$$

References

- (AbM) Abramowitz M., Stegun A., Handbook of mathematical functions, Dover Publications Inc., New York, 1970
- (AhJ) Ahner J. F., A scattering trinity: the reproducing kernel, null-field equations and modified Green's functions, Q. J. Mech. Appl. Math. Vol. 39, No. 1, 153-162, 1986
- (AlF) Almgren F. J., Plateau's Problem, An Invitation to Varifold Geometry, American Mathematical Society, New York, 2001
- (AnE) Anderson E., The Problem of Time, Springer, Cambridge, UK, 2017
- (AnJ) Annett J. F., Superconductivity, Superfluids and condensates, Oxford University Press, Oxford, 2004
- (AnM) Anderson M. T., Geometrization of 3-manifolds via the Ricci flow, Notices Amer. Math. Soc. 51, (2004) 184-193
- (ApT) Apostol T. M., Introduction to Analytic Number Theory, Springer Verlag, 2000
- (ArA) Arthurs A. M., Complementary Variational Principles, Clarendon Press, Oxford, 1970
- (ArE) Artin E., The Gamma Function, Holt, Rinehart and Winston, New York, Chicago, San Francisco, Toronto, London, 1964
- (ArN) Arcozzi N., Li X., Riesz transforms on spheres, Mathematical Research Letters, 4, 401-412, 1997
- (AxC) Axler C. On a Family of Functions Defined Over Sums of Primes, Journal of Integer Sequences, Vol. 22 (2019), Article 19.5.7, christian.axler@hhu.de
- (AzA) Aziz A. K., Kellog R. B., Finite Element Analysis of Scattering Problem, Math. Comp., Vol. 37, No 156 (1981) 261-272
- (AzT) Azizov T. Y., Ginsburg Y. P., Langer H., On Krein's papers in the theory of spaces with an indefinite metric, Ukrainian Mathematical Journal, Vol. 46, No 1-2, 1994, 3-14
- (AzT1) Azizov, T. Y., Iokhvidov, I. S., Dawson, E. R., Linear Operators in Spaces With an Indefinite Metric, Wiley, Chichester, New York, 1989
- (BaB) Bagchi B., On Nyman, Beurling and Baez-Duarte's Hilbert space reformulation of the Riemann Hypothesis, Indian Statistical Institute, Bangalore Centre, (2005), www.isibang.ac.in
- (BaÁ) Baricz Á., Mills' ratio: Monotonicity patterns and functional inequalities, J. Math. Anal. Appl. 340, 1362-1370, 2008
- (BaÀ) Baricz Á., Pogány T. K., Inequalities for the one-dimensional analogous of the Coulomb potential, Acta Polytechnica Hungarica, Vol. 10, No. 7, 53-67, 2013
- (BaR) Barnard R. W., Gordy M., Richards K. C., A note on Turán type and mean inequalities for the Kummer function, J. Math. Anal. Appl. 349 (1), 259-263, 2009
- (BeA) Besse A., L., Einstein Manifolds, Springer-Verlag, Berlin, Heidelberg, 1987
- (BeB) Berndt B. C., Ramanujan's Notebooks, Part I, Springer Verlag, New York, Berlin, Heidelberg, Tokyo, 1985
- (BeB1) Berndt B. C., Number Theory in the Spirit of Ramanujan, AMS, Providence, Rhode Island, 2006
- (BeG) Besson G., The geometrization conjecture after R. Hamilton and G. Perelman, Rend. Sem. Mat. Pol. Torino, Vol. 65, 4, 2007, pp. 397-411

- (BeI) Belogrivov, I. I., On Transcendence and algebraic independence of the values of Kummer's functions, Translated from Sibirskii Matematicheskii Zhurnal, Vol. 12, No 5, 1971, 961-982
- (BeL) Bel L., Introduction d'un tenseur du quatrième ordre, C. R. Acad. Sci. Paris, 247, 1094-1096, 1959
- (BiD) Bleecker D., Gauge Theory and Variational Principles, Dover Publications, Inc., Mineola, New York, 1981
- (BiN) Bingham N. H., Goldie C. M., Teugels J. L., Regular variation, University Press, Cambridge, 1989
- (BiN1) Bingham N. H., Szegő's theorem and its probabilistic descendants, Probability Surveys, Vol. 9, 287-324 2012
- (BiP) Biane P., Pitman J., Yor M., Probability laws related to the Jacobi theta and Riemann Zeta functions, and Brownian excursion, Amer. Math. soc., Vol 38, No 4, 435-465, 2001
- (BoD) Bohm D., Wholeness and the Implicate Order, Routledge & Kegan Paul, London, 1980
- (BoJ) Bogner J., Indefinite Inner Product Spaces, Springer-Verlag, Berlin, Heidelberg, New York, 1974
- (BoM) Bonnet M., Boundary Integral Equations Methods for Solids and Fluids, John Wiley & Sons Ltd., Chichester, 1995
- (BrH) Bezis H., Asymptotic Behavior of Some Evolution Systems, In: Nonlinear Evolution Equations (M. C. Crandall ed.). Academic Press, New York, 141-154, 1978
- (BrK) Braun K., A new ground state energy model, www.quantum-gravitation.de
- (BrK1) Braun K., An alternative Schroedinger (Calderon) momentum operator enabling a quantum gravity model
- (BrK2) Braun K., Global existence and uniqueness of 3D Navier-Stokes equations
- (BrK3) Braun K., Some remarkable Pseudo-Differential Operators of order -1, 0, 1
- (BrK4) Braun K., A Kummer function based Zeta function theory to prove the Riemann Hypothesis and the Goldbach conjecture
- (BrK5) An alternative trigonometric integral representation of the Zeta function on the critical line
- (BrK6) Braun K., A distributional Hilbert space framework to prove the Landau damping phenomenon
- (BrK7=BrK1) Braun K., An alternative Schroedinger (Calderón) momentum operator enabling a quantum gravity model
- (BrK8) Braun K., Comparison table, math. modelling frameworks for SMEP and GUT
- (BrK9) Braun K., Interior Error Estimates of the Ritz Method for Pseudo-Differential Equations, Jap. Journal of Applied Mathematics, 3, 1, 59-72, 1986
- (BrK10) Braun K., J. A. Nitsche's footprints to NSE problems, www.navier-stokes-equations.com
- (BrK related papers) www.navier-stokes-equations.com/author-s-papers
- (BuH) Buchholtz H., The Confluent Hypergeometric Function, Springer-Verlag, Berlin, Heidelberg, New York, 1969
- (BrR) Brent R. P., An asymptotic expansion inspired by Ramanujan, rpb@cslab.anu.edu.au
- (CaD) Cardon D., Convolution operators and zeros of entire functions, Proc. Amer. Math. Soc., 130, 6 (2002) 1725-1734

- (CaH) Cao H.-D., Zhu X.-P., Hamilton-Perelman's Proof of the Poincare Conjecture and the Geometrization Conjecture, arXiv:math/0612069v1, 2006
- (CaH1) Cao H. D., Chow B., Chu S. C., Yau S. T., Collected Papers on Ricci Flow, International Press 2003
- (CaJ) Cao J., DeTurck D., The Ricci Curvature with Rotational Symmetry, American Journal of Mathematics 116, (1994), 219-241
- (CeC) Cercignani C., Theory and application of the Boltzmann equation, Scottish Academic Press, Edinburgh, London, 1975
- (ChD) Christodoulou D., Klainerman, Asymptotic properties of linear field equations in Minkowski space, Comm. Pure Appl. Math. 43, 137-199, 1990
- (ChD1) Christodoulou D., Klainerman, The Global Nonlinear Stability of the Minkowski Space, Princeton University Press, New Jersey, 1993
- (ChF) Chen F., F., Introduction to Plasma Physics and Controlled Fusion, Volume I: Plasma Physics, Plenum Press, New York, 1984
- (ChH) Chen H., Evaluations of Some Variant Euler Sums, Journal of Integer Sequences, Vol. 9, (2006) 1-9
- (ChK) Chandrasekharan K., Elliptic Functions, Springer-Verlag, Berlin, Heidelberg, New York, Tokyo, 1985
- (ChK1) Chaudhury K. N., Unser M., On the Hilbert transform of wavelets, arXiv:1107.4619v1
- (ChJ) Choi J., Srivastava H. M., The Clausen function and its related integrals, Thai J. Math., Vol 12, No 2, 251-264, 2014
- (ChO) Christensen O., An Introduction to Frames and Riesz Bases, Applied and Numerical Harmonic Analysis, Second Edition, Birkhäuser, 2015
- (CiI) Ciufolini I., Wheeler J. A., Gravitation and Inertia, Princeton University Press, Princeton, New Jersey, 1995
- (CoF) Coffey M. W., Polygamma theory, Li/Kneiper constants, and validity of the Riemann Hypothesis, <http://arxiv.org>
- (CoG) Cohen G., Lin M., The One-sided Ergodic Hilbert Transform of Normal Contractions, Operator Theory: Advances and Applications, Vol. 197, 77-98 c 2009 Birkhäuser Verlag Basel, Switzerland
- (CoP) Constantin P., Lax P. D., Majda A., A simple one-dimensional model for the three-dimensional vorticity model, Communications on Pure and Applied Mathematics, Vol. XXXVIII, 715-724, 1985
- (CoR) Courant R., Hilbert D., Methods of Mathematical Physics Volume II, J. Wiley & Sons, New York, 1989
- (DeJ) Dereziński J., Richard S., On Radial Schrödinger Operators with a Coulomb Potential, Ann. Henri Poincaré 19 (2018), 2869-2917
- (DeJ) Derbyshire J., Prime Obsession, Joseph Henry Press, Washington, D. C., 2003
- (DeR) Dendy R. O., Plasma Dynamics, Oxford Science Publications, Oxford, 1990
- (DrM) Dritschel M. A., Rovnyak, J., Operators on indefinite inner product spaces
- (DuR) Duffin R. J., Schaeffer A. C., A Class of Nonharmonic Fourier Series, Transactions of the American Mathematical Society, Vol. 72, No. 2, 1952, pp. 341-366
- (EbP) Ebenfelt P., Khavinson D., Shapiro H. S., An inverse problem for the double layer potential, Computational Methods and Function Theory, Vol. 1, No. 2, 387-401, 2001

- (EdH) Edwards Riemann's Zeta Function, Dover Publications, Inc., Mineola, New York, 1974
- (EhP) Ehrlich P., Contemporary infinitesimalist theories of continua and their late 19th- and early 20th-century forerunners, arXiv.org 180.03345, Dec 2018
- (EiA) Einstein A., Grundzüge der Relativitätstheorie, Vieweg & Sohn, Braunschweig, Wiesbaden, 1992
- (EiA1) Einstein A., Äther und Relativitätstheorie, Julius Springer, Berlin, 1920
- (EiA2) Einstein A., Podolsky B., Rosen N., Can Quantum-Mechanical Description of Physical Reality Be Considered Complete? Physical Review, Vol. 47, 1935
- (EIL) Elaissaoui L., Guennoun Z. El-Abidine, Relating log-tangent integrals with the Riemann zeta function, arXiv, May 2018
- (EIL1) Elaissaoui L., Guennoun Z. El-Abidine, Evaluation of log-tangent integrals by series involving $\zeta(2n+1)$, arXiv, May 2017
- (EsG) Eskin G. I., Boundary Value Problems for Elliptic Pseudodifferential Equations, Amer. Math. Soc., Providence, Rhode Island, 1981
- (EsO) Esinosa O., Moll V., On some definite integrals involving the Hurwitz zeta function, Part 2, The Ramanujan Journal, 6, p. 449-468, 2002
- (EsR) Estrada R., Kanwal R. P., Asymptotic Analysis: A Distributional Approach, Birkhäuser, Boston, Basel, Berlin, 1994
- (EyG) Eyink G. L., Stochastic Line-Motion and Stochastic Conservation Laws for Non-Ideal Hydrodynamic Models. I. Incompressible Fluids and Isotropic Transport Coefficients, arXiv:0812.0153v1, 30 Nov 2008
- (FaK) Fan K., Invariant subspaces of certain linear operators, Bull. Amer. Math. Soc. 69 (1963), no. 6, 773-777
- (FaM) Farge M., Schneider K., Wavelets: application to turbulence, University Warnick, lectures, 2005
- (FaM1) Farge M., Schneider K., Wavelet transforms and their applications to MHD and plasma turbulence: a review, arXiv:1508.05650v1, 2015
- (FeR) Feynman R. P., Quantum Electrodynamics, Benjamin/Cummings Publishing Company, Menlo Park, California, 1961
- (FID) Fleisch D., A Student's Guide to Maxwell's Equations, Cambridge University Press, 2008
- (GaA) Ganchev A. H., Greenberg W., van der Mee C. V. M., A class of linear kinetic equations in Krein space setting, Integral Equations and Operator Theory, Vol. 11, 518-535, 1988
- (GaG) Galdi G. P., The Navier-Stokes Equations: A Mathematical analysis, Birkhäuser Verlag, Monographs in Mathematics, ISBN 978-3-0348-0484-4
- (GaL) Garding L., Some points of analysis and their history, Amer. Math. Soc., Vol. 11, Providence Rhode Island, 1919
- (GaW) Gautschi W., Waldvogel J., Computing the Hilbert Transform of the Generalized Laguerre and Hermite Weight Functions, BIT Numerical Mathematics, Vol 41, Issue 3, pp. 490-503, 2001
- (GiY) Giga Y., Weak and strong solutions of the Navier-Stokes initial value problem, Publ. RIMS, Kyoto Univ. 19 (1983) 887-910
- (GöK) Gödel, K., An Example of a New Type of Cosmological Solutions of Einstein's Field Equations of Gravitation, Review of Modern Physics, Vol. 21, No. 3, 1949

- (GrI) Gradshteyn I. S., Ryzhik I. M., Table of integrals series and products, Academic Press, New York, San Francisco, London, 1965
- (GrJ) Graves J. C., The conceptual foundations of contemporary relativity theory, MIT Press, Cambridge, Massachusetts, 1971
- (GrS) Graham S. W., Van der Corput's Method of Exponential Sums, Cambridge University Press, Cambridge, 1991
- (GuR) Gundersen R. M., Linearized Analysis of One-Dimensional Magnetohydrodynamic Flows, Springer Tracts in Natural Philosophy, Vol 1, Berlin, Göttingen, Heidelberg, New York, 1964
- (HaE) Haidemenakis E. D., Physics of Solids in Intense Magnetic Fields, Plenum Press, New York, 1969
- (HaG) Hardy G. H., Riesz M., The general theory of Dirichlet's series, Cambridge University Press, Cambridge, 1915
- (HaH) Halberstam H., Richert H.-E., Sieve Methods, Academic Press, London, New-York, San Francisco 1974
- (HaJ) Havil J., Gamma, exploring euler's constant, Princeton University Press, Princeton and Oxford, 2003
- (HaJ1) Hartle J. B., Hawking S. W., Wave function of the Universe, Physical Review, D, Vol. 28, No. 12, 1983
- (HaP) Halmos P. R., Lectures on Ergodic Theory, Martino Publishing, Mansfield Centre, CT, 2013
- (HaR) Hamilton R. S., Non-singular solutions of the Ricci flow on three-manifolds, Com. Anal. and Geometry, Vol. 7, No. 4, pp. 695-729, 1999
- (HaR1) Hamilton R. S., Three manifolds with positive Ricci curvature, Jour. Diff. Geom. 17, pp. 255-306, 1982
- (HaS) Hawking S. W., Penrose R., The Singularities of Gravitational Collapse and Cosmology, The Royal Society, Vol. 314, Issue 1519, 1970
- (HaS1) Hawking S. W., Particle Creation by Black Holes, Commun. Math. Phys. 43, 199-220, 1975
- (HeJ) Heywood J. G., Walsh O. D., A counter-example concerning the pressure in the Navier-Stokes equations, as $t \rightarrow 0^+$, Pacific Journal of Mathematics, Vol. 164, No. 2, 351-359, 1994
- (HeM) Heidegger M., Holzwege, Vittorio Klostermann, Frankfurt a. M., 2003
- (HeW) Heisenberg W., Physikalische Prinzipien der Quantentheorie, Wissenschaftsverlag, Mannheim, Wien, Zürich, 1991
- (HiF) Hirzebruch F., Scharlau W., Einführung in die Funktionalanalysis, Spektrum Akademischer Verlag, Heidelberg, Berlin, Oxford, 1996
- (HiJ) Higgins J. R., Five short stories about the cardinal series, Bulletin of the American Mathematical Society, Vol. 12, No. 1, 1985, pp. 45-89
- (HiP) Higgs P. W., Spontaneous Symmetry Breakdown without Massless Bosons, Physical Review, Vol. 145, No 4, p. 1156-1162, 1966
- (HoE) Hopf E., Ergodentheorie, Verlag von Julius Springer, Berlin, 1937
- (HoA) Horvath A. G., Semi-indefinite-inner-product and generalized Minkowski spaces, arXiv
- (HoM) Holschneider M., Wavelets, An Analysis Tool, Clarendon Press, Oxford, 1995
- (HuA) Hurwitz A., Über einen Satz des Herrn Kakeya, Zürich, 1913

- (IvV) Ivakhnenko, V. I., Smirnow Yu. G., Tyrtshnikov E. E., The electric field integral equation: theory and algorithms, Inst. Numer. Math. Russian of Academy Sciences, Moscow, Russia
- (JoF) John F., Formation of singularities in elastic waves, Lecture Notes in Phys., Springer-Verlag, 190-214, 1984
- (KaD) Karp D., Sitnik S. M., Log-convexity and log-concavity of hypergeometric-like functions, J. Math. Anal. Appl. 364, 384-394, 2010
- (KaD1) Kazanas D., Cosmological Inflation: A Personal Perspective, Astrophys. Space Sci. Proc., (2009) 485-496
- (KaM) Kaku M., Introduction to Superstrings and M-Theory, Springer-Verlag, New York, Inc., 1988
- (KaM1) Kac M., Probability methods in some problems of analysis and number theory, Bull. Am. Math. Soc., 55, 641-655, (1949)
- (KeL) Keiper J. B., Power series expansions of Riemann's Zeta function, Math. Comp. Vol 58, No 198, (1992) 765-773
- (KiA) Kirsch A., Hettlich F., The mathematical theory of time-harmonic Maxwell's equations, expansion-, integral-, and variational methods, Springer-Verlag, Heidelberg, New York, Dordrecht, London, 2015
- (KiA1) Kiselev A. P., Relatively Undistorted Cylindrical Waves Depending on Three Spatial Variables, Mathematical Notes, Vol. 79, No. 4, 587-588, 2006
- (KIB) Kleiner B., Lott J., Notes on Perelman's papers, Mathematics ArXiv
- (KiH) Kim H., Origin of the Universe: A Hint from Eddington-inspired Born-Infeld gravity, Journal of the Korean Physical Society, Vol. 65, No. 6, pp. 840-845, 2014
- (KIS) Klainermann S., Rodnianski, Regularity and geometric properties of solutions of the Einstein-Vacuum equations, Journées équations aux dérivées partielles, No. 14, p. 14, 2002
- (KIS1) Klainerman S., Nicolò, The Evolution Problem in General Relativity, Birkhäuser, Boston, Basel, Berlin, 1950
- (KIS2) Klainerman S., Remarks on the global Sobolev inequalities in Minkowski space, Comm. Pure. Appl. Math., 40, 111-117, 1987
- (KIS3) Klainerman S., Uniform decay estimates and the Lorentz invariance of the classical wave equation, Comm. Pure Appl. Math., 38, 321-332, 1985
- (KnA) Kneser A., Das Prinzip der kleinsten Wirkung von Leibniz bis zur Gegenwart, B. G. Teubner, Leipzig, Berlin, 1928
- (KoA) Kolmogoroff A., Une contribution à l'étude de la convergence des séries de Fourier, Fund. Math. Vol. 5, 484-489
- (KoB) Koopman B. O., Hamiltonian Systems and Transformations in Hilbert Spaces, PNAS, 1931, 17 (5), pp. 315-318
- (KoB1) Koopman B. O., v. Neumann J., Dynamical Systems of continuous spectra, PNAS, 1932, 18 (3), pp. 255-263
- (KoG) Kozma G., Lev N., Exponential Riesz bases, discrepancy of irrational rotations and BMO, J. Fourier Anal. Appl. 17, 2011, pp 879-898
- (KoH) Koch H., Tataru D., Well-posedness for the Navier-Stokes equations, Adv. Math., Vol 157, No 1, 22-35, 2001
- (KoJ) Korevaar J., Distributional Wiener-Ikehara theorem and twin primes, Indag. Mathem. N. S., 16, 37-49, 2005

- (KrA) Krall A. M., Spectral Analysis for the Generalized Hermite Polynomials, Trans. American Mathematical Society, Vol. 344, No. 1 (1994) pp. 155-172
- (KrR) Kress R. Linear Integral Equations, Springer-Verlag, Berlin, Heidelberg, New York, London, Paris, Tokyo, Hong Kong, 1941
- (KrR1) Kraußhar R. S., Malonek H. R., A characterization of conformal mappings in R^4 by a formal differentiability condition, Bulletin de la Societè Royale de Liège, Vol. 70, Vol. 1, 35-49, 2001
- (LaC) Lanczos C., The variational principles of mechanics, Dover Publications Inc., New York, 1970
- (LaC1) Langenhof C. E., Bounds on the norm of a solution of a general differential equation, Proc. Amer. Math. Soc., 8, 615-616, 1960
- (LaE) Landau E., Die Lehre von der Verteilung der Primzahlen Vol 1, Teubner Verlag, Leipzig Berlin, 1909
- (LaE1) Landau E., Ueber die zahlentheoretische Function $\varphi(n)$ und ihre Beziehung zum Goldbachschen Satz, Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse, Vol 1900, p. 177-186, 1900
- (LaE2) Landau E., Die Lehre von der Verteilung der Primzahlen Vol 2, Teubner Verlag, Leipzig Berlin, 1909
- (LaE3) Landau E., Über eine trigonometrische Summe, Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse, Vol 1928, p. 21-24, 1928
- (LaE4) Landau E. Vorlesungen über Zahlentheorie, Erster Band, zweiter Teil, Chelsea Publishing Company, New York, 1955
- (LaE5) Landau E., Die Goldbachsche Vermutung und der Schnirelmannsche Satz, Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Math.-Phys. Klasse, 255-276, 1930
- (LaE6) Landau E., Über die Fareyreihe und die Riemannsche Vermutung, Göttinger Nachrichten (1932), 347-352
- (LaG) Lachaud G., Spectral analysis and the Riemann hypothesis, J. Comp. Appl. Math. 160, pp. 175190, 2003
- (LaJ) An Elementary Problem Equivalent to the Riemann Hypothesis, <https://arxiv.org>
- (LaG1) Lagarias J. C., The Schrödinger Operator with Morse Potential on the Right Half Line, arXiv:0712.3238v6
- (LeN) Lebedev N. N., Special Functions and their Applications, translated by R. A. Silverman, Prentice-Hall, Inc., Englewood Cliffs, New York, 1965
- (LeN1) Lerner N., A note on the Oseen kernels, in Advances in phase space analysis of partial differential equations, Siena, pp. 161-170, 2007
- (LeP) LeFloch P. G., Ma Y., The global nonlinear stability of Minkowski space, arXiv: 1712.10045v1, 28 Dec 2017
- (LiI) Lifanov I. K., Poltavskii L. N., Vainikko G. M., Hypersingular Integral Equations and their Applications, Chapman & Hall/CRC, Boca Raton, London, New York, Washington, D. C., 2004
- (LiI1) Lifanov I. K., Nenashaev A. S., Generalized Functions on Hilbert Spaces, Singular Integral Equations, and Problems of Aerodynamics and Electrodynamics, Differential Equations, Vol. 43, No. 6, pp. 862-872, 2007
- (LiJ) Linnik J. V., The dispersion method in binary additive problems, American Mathematical Society, Providence, Rhode Island, 1963

- (LiP) Lions P. L., On Boltzmann and Landau equations, *Phil. Trans. R. Soc. Lond. A*, 346, 191-204, 1994
- (LiP1) Lions P. L., Compactness in Boltzmann's equation via Fourier integral operators and applications. III, *J. Math. Kyoto Univ.*, 34-3, 539-584, 1994
- (LiX) Li Xian-Jin, The Positivity of a Sequence of Numbers and the Riemann Hypothesis, *Journal of Number Theory*, 65, 325-333 (1997)
- (LoA) Lifanov I. K., Poltavskii L. N., Vainikko G. M., *Hypersingular Integral Equations and Their Applications*, Chapman & Hall/CRC, Boca Raton, London, New York, Washington, D. C. 2004
- (LoJ) Long J. W., *The Grad Student's Guide to Kant's Critique of Pure Reason*, iUniverse, Bloomington, 2016
- (LuL) Lusternik L. A., Sobolev V. J., *Elements of Functional Analysis*, A Halsted Press book, Hindustan Publishing Corp. Delhi, 1961
- (MaJ) Mashreghi, J., Hilbert transform of $\log(\text{abs}(f))$, *Proc. Amer. Math. Soc.*, Vol 130, No 3, p. 683-688, 2001
- (MaJ1) Marsden J. E., Hughes T. J. R., *Mathematical foundations of elasticity*, Dover Publications Inc., New York, 1983
- (MeJ) Meiklejohn J. M. D., *The Critique of Pure Reason*, By Immanuel Kant, Translated by J. M. D. Meiklejohn, ISBN-13 978-1977857477, ISBN-10: 1977857477
- (MeY) Meyer Y., Coifman R., *Wavelets, Calderón-Zygmund and multilinear operators*, Cambridge University Press, Cambridge, 1997
- (MiJ) Milnor J., *Morse Theory*, *Annals of Mathematical Studies*, Number 51, Princeton University Press, Princeton, 963
- (MiM) Mikolas M., Farey series and their connection with the prime number problem I. *Acta Univ. Szeged. Sect. Sci. Math.* 13 (1949), 93-117.
- (MiM1) M. Mikolas M., Farey series and their connection with the prime number problem II. *Acta Univ. Szeged. Sect. Sci. Math.* 14 (1951), 5-21.
- (MiT) Mikosch T., *Regular Variation, Subexponentiality and Their Application in Probability Theory*, University of Groningen
- (MoC) Mouhot C., Villani C., On Landau damping, *Acta Mathematica*, Vol. 207, Issue 1, p. 29-201, 2011
- (MoJ) Morgan J. W., Tian G., Ricci Flow and the Poincare Conjecture, *Mathematics ArXiv*
- (MoM) Morse M., Functional Topology and Abstract Variational Theory, *Proc. N. A. S.*, 326-330, 1938
- (NaA) Nakamura A., **Basis properties and complements of complex exponential systems**, *Hokkaido Mathematical Journal* Vol. 36 (2007) p. 193-206
- (NaC) Nasim C. On the summation formula of Voronoi, *Trans. American Math. Soc.* 163, 35-45, 1972
- (NaP) Naselsky P. D., Novikov D. I., Noyikov I. D., *The Physics of the Cosmic Microwave Background*, Cambridge University Press, Cambridge, New York, Melbourne, Madrid, Cape Town, Singapore, Sao Paulo, 2006
- (NaS) Nag S., Sullivan D., Teichmüller theory and the universal period mapping via quantum calculus and the H space on the circle, *Osaka J. Math.*, 32, 1-34, 1995
- (NeD) Neuenschwander D. E., *Emmy Noether's Wonderful Theorem*, The John Hopkins University Press, Baltimore, 2011
- (NiJ) Nitsche J. A., lecture notes, approximation theory in Hilbert scales

- (NiJ1) Nitsche J. A., lecture notes, approximation theory in Hilbert scales, extensions and generalizations
- (NiJ2) Nitsche J. A., Direct Proofs of Some Unusual Shift-Theorems, *Anal. Math. Appl.*, Gauthier-Villars, Montrouge, 1988, pp. 383–400
- (NiJ3) Nitsche J. A., Free boundary problems for Stokes' s flows and finite element methods, *Equadiff* 6, 2006, pp. 327-332
- (NiN) Nielsen N., *Die Gammafunktion*, Chelsea Publishing Company, Bronx, New York, 1965
- (ObF) Oberhettinger, *Tables of Mellin Transforms*, Springer-Verlag, Berlin, Heidelberg, New York, 1974
- (OIF) Olver F. W. J., *Asymptotics and special functions*, Academic Press, Inc., Boston, San Diego, New York, London, Sydney, Tokyo, Toronto, 1974
- (OIF1) Olver F. W. J., Lozier D. W., Boisvert R. F., Clark C. W., *NIST Handbook of Mathematical Functions*
- (OIJ) Olsen J.-F., Local Properties of Hilbert spaces of Dirichlet series, *Journal of Functional Analysis* 262, 2011, pp. 2669-2696
- (OIR) Oloff R., *Geometrie der Raumzeit*, Vieweg & Sohn, Braunschweig/Wiesbaden, 1999
- (OsK) Oskolkov K. I., Chakhkiev M. A., On Riemann „Nondifferentiable“ Function and the Schrödinger Equation, *Proc. Steklov Institute of Mathematics*, Vol. 269, 2010, pp. 186-196
- (OsH) Ostmann H.-H., *Additive Zahlentheorie, erster Teil*, Springer-Verlag, Berlin, Göttingen, Heidelberg, 1956
- (PeB) Petersen B. E., *Introduction the the Fourier transform and Pseudo-Differential operators*, Pitman Advanced Publishing Program, Boston, London, Melbourne, 1983
- (PeM) Perel M., Gorodnitskiy E., Representations of solutions of the wave equation based on relativistic wavelet, *arXiv:1205.3461v1*, 2012
- (PeO) Perron O., *Die Lehre von den Kettenbrüchen, Volumes 1-2*, Nabu Public Domain Reprint, copyright 1918 by B. G. Teubner in Leipzig
- (PeR) Penrose R., *Cycles of Time*, Vintage, London, 2011
- (PeR1) Peralta-Fabi, R., An integral representation of the Navier-Stokes Equation-I, *Revista Mexicana de Fisica*, Vol 31, No 1, 57-67, 1984
- (PeR2) Penrose R., *Structure of space-time*, *Batelle Rencontre*, C. M. DeWitt and J. M. Wheeler, 1967
- (PeR3) Penrose R., Zero rest mass fields including gravitation: asymptotic behaviours, *Proc. Toy. Soc. Lond.*, A284, 159-203, 1962
- (PeR4) Penrose R., *The Emperor's New Mind: Concerning Computers, Minds, and the Laws of Physics*, Oxford Univ. Press, 1989
- (PeR5) Penrose R., Rindler W., *Spinors and Space-Time*, Cambridge University Press, Cambridge, 1984
- (PhR) Phillips R., Dissipative operators and hyperbolic systems of partial differential equations, *Trans. Amer. Math. Soc.* 90 (1959), 193-254
- (PiS) Pilipovic S., Stankovic B., Tauberian Theorems for Integral Transforms of Distributions, *Acta Math. Hungar.* 74, (1-2) (1997), 135-153
- (PiS1) Pilipovic S., Stankovic B., Wiener Tauberian theorems for distributions, *J. London Math. Soc.* 47 (1993), 507-515

- (PIJ) J. Plemelj, Potentialtheoretische Untersuchungen, B.G. Teubner, Leipzig, 1911
- (PoE) Postnikov E. B., Singh V. K. Continuous wavelet transform with the Shannon wavelet from the point of view of hyperbolic partial differential equations, arXiv:1551.03082
- (PoG) Polya G., Über Nullstellen gewisser ganzer Funktionen, Math. Z. 2 (1918) 352-383
- (PoG1) Polya G., Über eine neue Weise bestimmte Integrale in der analytischen Zahlentheorie zu gebrauchen, Göttinger Nachr. (1917) 149-159
- (PoG2) Polya G. Über die algebraisch-funktionentheoretischen Untersuchungen von J. L. W. V. Jensen, Det Kgl. Danske Videnskabernes Selskab., Mathematisk-fysiske Meddelelser. VII, 17, 1927
- (PoG3) Polya G., Über Potenzreihen mit ganzzahligen Koeffizienten, Math. Ann. 77, 1916, 497-513
- (PoG4) Polya G., Arithmetische Eigenschaften der Reihenentwicklung rationaler Funktionen, J. Reine und Angewandte Mathematik, 151, 1921, 1-31
- (PoD) Pollack D., Initial Data for the Cauchy Problem in General Relativity, General Relativity Spring School 2015, Junior Scientist Andrejewski Days, March 22nd to April 4th, 2015, Brandenburg an der Havel, Germany
- (PoP) Poluyan P., Non-standard analysis of non-classical motion; do the hyperreal numbers exist in the quantum-relative universe?
- (PrK) Prachar K., Primzahlverteilung, Springer-Verlag, Berlin, Göttingen, Heidelberg, 1957
- (ReR) Redheffer R., Two consequences of the Beurling-Malliavin theory, Proc. Amer. Math. Soc., Volume 36, No 1, 1972, pp. 116-122
- (RiB) Riemann B., Ueber die Darstellbarkeit einer Function durch eine trigonometrische Reihe, Abhandlungen der Königlichen Gesellschaft der Wissenschaften zu Göttingen, transcribed by D. R. Wilkins, 2000
- (RoC) Rovelli C., Quantum Gravity, Cambridge University Press, Cambridge, 2004
- (RoC1) Rovelli C., The Order of Time, Penguin Random House, 2018
- (RoC2) Rovelli C., Reality is not what it seems, Penguin books, 2017
- (RoC3) Rovelli C., Seven brief lessons on physics, Penguin Books, 2016
- (RoJ) Roberts J. T., Leibniz on Force and Absolute Motion, Philosophy of Science, Vol 70, No 3, pp. 553-573, 2003
- (RuB) Russel B., The Philosophy of Leibniz, Routledge, London, New York, paperback edition, 1992
- (RuC) Runge C., Über eine Analogie der Cauchy-Riemannschen Differentialgleichungen in drei Dimensionen, Nachr. v. d. Gesellschaft d. Wissenschaften zu Göttingen, Math-Phys. Klasse Vol 1992, 129-136, 1992
- (RuM) Ruskai M. B., Werner E., Study of a Class of Regularizations of $1/|x|$ using Gaussian Integrals, arXiv:math/990212v2
- (RyG) Rybicki, G. B., Dawson's integral and the sampling theorem, Computers in Physics, 3, 85-87, 1989
- (ScE) Schrödinger E., Statistical Thermodynamics, Dover Publications, Inc., New York, 1989
- (ScE1) Schrödinger E., My View of the World, Ox Bow Press, Woodbridge, Connecticut, 1961
- (ScE2) Schrödinger E., *What is Life?* and *Mind and Matter*, Cambridge University Press, Cambridge, 1967

- (ScP) Scott P., The Geometries of 3-Manifolds, Bull. London Math. Soc., 15 (1983), 401-487
- (SeA) Sedletsii A. M., Asymptotics of the Zeros of Degenerated Hypergeometric Functions, Mathematical Notes, Vol. 82, No. 2, 229-237, 2007
- (SeE) Seneta E., Regularly Varying Functions, Lecture Notes in Math., 508, Springer Verlag, Berlin, 1976
- (SeH) Seifert H., Threlfall W., Variationsrechnung im Grossen, Chelsea Publishing Company, New York, 1951
- (SeK) Seip K., On the Connection between Exponential Bases and Certain Related Sequences in $L_2(-\pi, \pi)$, Journal of Functional Analysis 130, 1995, pp. 131-160
- (SeJ) Serrin J., Mathematical Principles of Classical Fluid Mechanics
- (ShF) Shu F. H., Gas Dynamics, Vol II, University Science Books, Sausalito, California, 1992
- (ShM) Scheel M. A., Thorne K. S., Geodynamics, The Nonlinear Dynamics of Curved Spacetime
- (ShM1) Shimoji M., Complementary variational formulation of Maxwell's equations in power series form
- (SiT) Sideris T., Formation of singularities in 3-D compressible fluids, Comm. Math. Phys., 101, 47-485, 1985
- (SmL) Smolin L., Time reborn, Houghton Mifflin Harcourt, New York, 2013
- (SmL1) Smith L. P., Quantum Effects in the Interaction of Electrons With High Frequency Fields and the Transition to Classical Theory, Phys. Rev. 69 (1946) 195
- (SoH) Sohr H., The Navier-Stokes Equations, An Elementary Functional Analytical Approach, Birkhäuser Verlag, Basel, Boston, Berlin, 2001
- (SoP) Sobolevskii P. E., On non-stationary equations of hydrodynamics for viscous fluid. Dokl. Akad. Nauk SSSR 128 (1959) 45-48 (in Russian)
- (StE) Stein E. M., Conjugate harmonic functions in several variables
- (StE1) Stein E. M., Harmonic Analysis, Real-Variable Methods, Orthogonality, and Oscillatory Integrals, Princeton University Press, Princeton, New Jersey, 1993
- (SzG) Szegő, G., Orthogonal Polynomials, American Mathematical Society, Providence, Rhode Island, 2003
- (TaM) Tajmar M., de Mantos C. J., Coupling of Electromagnetism and Gravitation in the Weak Field Approximation, <https://arxiv.org/>
- (ThW) Thurston W. P., Three Dimensional Manifolds, Kleinian Groups and Hyperbolic Geometry, Bulletin American Mathematical Society, Vol 6, No 3, 1982
- (TiE) Titchmarsh E. C., The theory of the Riemann Zeta-function, Clarendon Press, London, Oxford, 1986
- (TrH) Treder H.-J., Singularitäten in der Allgemeinen Relativitätstheorie, Astron. Nachr. Bd. 301, H. 1, 9-12, 1980
- (TsB) Al'Tsuler B. L., Integral form of the Einstein equations and a covariant formulation of the Mach's principle, Soviet Physics Jetp, Vol. 24, No. 4, 1967
- (VaM) Vainberg M. M., Variational Methods for the Study of Nonlinear Operators, Holden-Day, Inc., San Francisco, London, Amsterdam, 1964
- (VeG) Veneziano G., A simple/short introduction to pre-big-bang physics/cosmology, in „Erice 1997, Highlights of subnuclear physics“ 364-380, talk given at conference: C97-08-26.2 p. 364-380

- (VeW) Velte W., Direkte Methoden der Variationsrechnung, B. G. Teubner, Stuttgart, 1976
- (ViI) Vinogradov I. M., The Method of Trigonometrical Sums in the Theory of Numbers, Dover Publications Inc., Mineola, New York 2004
- (ViI1) Vinogradov, I. M., Representation of an odd number as the sum of three primes, Dokl. Akad. Nauk SSSR 15, 291-294 (1937)
- (ViJ) Vindas J., Estrada R., A quick distributional way to the prime number theorem, Indag. Mathem., N.S. 20 (1) (2009) 159-165
- (ViJ1) Vindas J., Local behavior of distributions and applications, Dissertation, Department of Mathematics, Louisiana State University, 2009
- (ViJ2) Vindas J., Introduction to Tauberian theory, a distributional approach, <https://cage.ugent.be>
- (ViM) Villarino M. B., Ramanujan's Harmonic Number Expansion Into Negative Powers of a Triangular Number, Journal of Inequalities in pure and applied mathematics, Vol. 9, No. 3 (2008), Art. 89, 12 pp.
- (ViV) Vladimirov V. S., Drozzinov Yu. N., Zavalov B. I., Tauberian Theorems for Generalized Functions, Kluwer Academic Publishers, Dordrecht, Boston, London, 1988
- (WeD) Westra D. B., The Haar measure on SU(2), March 14, 2008
- (WeH*) Weyl H., Gravitation und Elektrizität, Sitzungsberichte Akademie der Wissenschaften Berlin, 1918, 465-48.
- (WeH) Weyl H., Space, Time, Matter, Cosimo Classics, New York, 2010
- (WeH1) Weyl H., Matter, structure of the world, principle of action, ... in (WeH) §34 ff.
- (WeH2) Weyl H., Was ist Materie? Verlag Julius Springer, Berlin, 1924
- (WeH3) Weyl H., Philosophy of Mathematics and Natural Science, Princeton University Press, Princeton and Oxford, 2009
- (WeH) Weyl H., Über die Gleichverteilung von Zahlen mod. Eins, Math. Ann., 77, 1914, 313-352
- (WeP) Werner P., Self-Adjoint Extension of the Laplace Operator with Respect to Electric and Magnetic Boundary Conditions, J. Math. Anal. Appl., 70, 1979, pp. 131-160
- (WeP1) Werner P., Spectral Properties of the Laplace Operator with respect to Electric and Magnetic Boundary Conditions, J. Math. Anal. Appl., 92, 1983, pp. 1-65
- (WhJ1) Whittaker J. M., Interpolatory Function Theory, Cambridge University Press, Cambridge, 1935
- (WhJ2) Whittaker J. M., The „Fourier“ Theory of Cardinal Functions, Proceedings of the Edinburgh Mathematical Society, Vol. 1, Issue 3, pp. 169-176, 1928
- (WhJ) Wheeler J. A., On the Nature of Quantum Geometrodynamics
- (WhJ1) Wheeler J. A., Awakening to the Natural State, Non-Duality Press, Salisbury, 2004
- (WhJ2) Wheeler J. A., At home in the universe, American Institute of Physics, Woodbury, 1996
- (WoJ) Wohlfart J., Werte hypergeometrischer Funktionen, Inventiones mathematicae, Vol. 92, Issue 1, 1988, 187-216
- (WoW) Wong W. Kant's Conception of Ether as a field in the Opus posthumum, Proc Eight Intern. Kant Congress, Marquette University Press, Vol. II, Memphis 1995

(YeR) Ye R., Ricci flow, Einstein metrics and space forms, Trans. Americ. Math. Soc., Vol. 338, No. 2, 1993

(YoR) Young R. M., An Introduction to Nonharmonic Fourier Series, Pure and Applied Mathematics, Academic Press Inc., New York, 1980

(ZeA) Zemanian A. H., Generalized Integral Transformations, Dover Publications, Inc. New York, 1968

(ZhB) Zhechev B., Hilbert Transform Relations

(ZyA) Zygmund A., Trigonometric series, Volume I & II, C